

# Mortality

**Ernesto F. L. Amaral**

April 07–09, 2020  
Population and Society (SOCI 312)

[www.ernestoamaral.com](http://www.ernestoamaral.com)



# Outline

- Introduction
- Measurement of mortality
  - Standardization
  - Life table
- Short history of mortality in the world
- Mortality trends and causes of death
- Mortality and longevity in the United States
- Infant mortality
- Future course of mortality
- Coronavirus pandemic




# Introduction

- The impact of mortality varies significantly according to social and demographic characteristics
  - Age
  - Sex
  - Race/ethnicity
  - Social class
  - Marital status
  - Area of residence
- Levels of development, medical conditions, and a host of other factors influence life expectancy

# Changes in causes of death

- There have been major changes over the historical record in the main causes of death
- People used to die mainly of infectious and parasitic diseases
- Today the major causes of death in developed countries are heart disease, cancer, and stroke
- Major causes of death are different in countries with high and low levels of life expectancy



**ATM** | **TEXAS A&M**  
UNIVERSITY.

# Measurement of mortality

- Quantification of mortality is central to demography
- Measurement of mortality dates back to John Graunt (1620–1674) and his analyses of the “Bills of Mortality”
- Mortality refers to the relative frequency of death in a population

# Two concepts of mortality

- Life span
  - Numerical age limit of human life
  - Maximum recorded age at death
  - 122 years and 164 days, lived by the Frenchwoman Jeanne Louise Calment
- Life expectancy
  - Average expected number of years of life to be lived by a particular population at a given time

# Crude death rate

- Crude death rate (*CDR*)

$$CDR = d / p * 1,000$$

- *d*: deaths in the year
- *p*: population at midyear

- Data for the United States for 2013

$$CDR = 2,596,993 / 316,497,531 * 1,000$$

$$CDR = 8.2$$

- World range of *CDR* in 2014

- United Arab Emirates (UAE) and Qatar = 1
- Lesotho = 21



# *CDR* and age composition

- When *CDRs* are compared among countries, differences are sometimes due to differences in age composition
  - Previous examples mean that there are 8 times more deaths per 1,000 people in the US than in the UAE
  - Why is the *CDR* of the US eight times higher than of the UAE?
  - The main reason is that the UAE is much younger in average age than is the U.S.
  - Younger people have lower death rates

# Young and old people

- Countries with
  - Large proportions of young people
  - Small proportions of old people
  - Usually have lower *CDRs*
- Countries with
  - Small proportions of young people
  - Large proportions of old people
  - Usually have higher *CDRs*



# Changing age structure

- If age structure has changed over time
  - CDRs should not be used to compare the death experiences of the same population at different points in time
- US *CDR* did not change much in 54 years
  - $CDR_{1960} = 9.5$  per 1,000
  - $CDR_{2014} = 8.2$  per 1,000
- *CDR* is not capable of capturing the reduction in mortality when the population becomes older
  - The US became older between 1960 and 2014
  - Median age: 29 in 1960 and 37 in 2014

# Crude rate

- *CDR* is a crude rate, because its denominator comprises the entire population
- However, population members are not equally at risk of experiencing death
- Risk of death varies by age, sex, race/ethnicity, socioeconomic status, and others
- Death rates vary considerably by age...



# Age-specific death rates


- Demographers use age-specific death rates as a more precise way to measure mortality
- Age-specific death rate ( ${}_nASDR_x$  or  ${}_nM_x$ )

$${}_nM_x = {}_nd_x / {}_np_x * 1,000$$

- ${}_nd_x$ : deaths to persons aged  $x$  to  $x+n$
- ${}_np_x$ : persons in the population who are aged  $x$  to  $x+n$
- $n$ : width of the age group
- $x$ : initial year of the age group
- For instance,  $ASDR$  for age group 15–19 is  ${}_5M_{15}$




# Age curve of mortality, US, 2007



Source: Minino, et al., 2009: 2.

Source: Poston, Bouvier 2017, p.169.



**ATM** | **TEXAS A&M**  
UNIVERSITY.

# Standardization

- ${}_nASDR_x$ , and not  $CDR$ , should be used to compare the mortality experiences of countries with different age compositions
  - We use **standardization** to take into account age composition when we compare death rates among different countries
  - We can compare crude death rates for different countries or years
  - We need to adjust for differences in age structure
  - We estimate age-adjusted death rates and apply to a standard population



# Mortality and fertility

- We cannot simply add up  ${}_nASDR_x$  and multiply by the width of the age interval
  - People die just once
- This makes sense for age-specific fertility rates ( ${}_nASFR_x$ ) and total fertility rates ( $TFR$ )

$$TFR = \sum({}_nASFR_x * n)$$

- $n$ : width of the age group
- Women can have more than one child

# Age standardization

- Young populations tend to have low *CDRs*
- Old populations tend to have high *CDRs*
- We estimate a variation of *CDR* that allows us to account for age composition when comparing death rates among different countries

$$CDR = \sum_n ASDR_x * ({}_nP_x / P) * 1,000$$

- $P$ : total population
- ${}_nP_x$ : population in age group  $x$
- ${}_nASDR_x$ : *ASDR* for people aged  $x$  to  $x+n$



# United States, 2006

Age group	Population	Prop. population	Deaths	Age-specific death rate (ASDR)
0-1	4,147,760	0.0139	27,126	0.00654
1-4	16,352,320	0.0548	4,742	0.00029
5-9	20,142,000	0.0675	2,820	0.00014
10-14	21,454,960	0.0719	3,862	0.00018
15-19	21,604,160	0.0724	13,827	0.00064
20-24	20,947,680	0.0702	19,062	0.00091
25-29	20,022,640	0.0671	18,020	0.00090
30-34	20,261,360	0.0679	21,477	0.00106
35-39	21,067,040	0.0706	32,233	0.00153
40-44	22,857,440	0.0766	52,801	0.00231
45-49	22,588,880	0.0757	77,028	0.00341
50-54	20,142,000	0.0675	99,300	0.00493
55-59	17,158,000	0.0575	127,312	0.00742
60-64	13,040,080	0.0437	149,961	0.01150
65-69	10,115,760	0.0339	180,061	0.01780
70-74	8,474,560	0.0284	234,830	0.02771
75-79	7,400,320	0.0248	321,914	0.04350
80-84	5,580,080	0.0187	388,262	0.06958
85-89	3,192,880	0.0107	353,005	0.11056
90-94	1,342,800	0.0045	234,681	0.17477
95-99	387,920	0.0013	107,283	0.27656
100+	89,520	0.0003	39,292	0.43892
<b>Total</b>	<b>298,400,000</b>	<b>0.9999</b>	<b>2,508,899</b>	<b>1.20116</b>

CDR per 1,000

8.41



# Venezuela, 2006

Age group	Population	Prop. population	Deaths	Age-specific death rate (ASDR)
0-1	487,056	0.0219	7,905	0.01623
1-4	1,779,200	0.0800	1,139	0.00064
5-9	2,308,512	0.1038	739	0.00032
10-14	2,275,152	0.1023	933	0.00041
15-19	2,250,688	0.1012	3,286	0.00146
20-24	2,057,200	0.0925	4,896	0.00238
25-29	1,872,608	0.0842	4,138	0.00221
30-34	1,614,624	0.0726	3,278	0.00203
35-39	1,530,112	0.0688	3,290	0.00215
40-44	1,401,120	0.0630	4,049	0.00289
45-49	1,145,360	0.0515	4,707	0.00411
50-54	971,888	0.0437	5,520	0.00568
55-59	769,504	0.0346	5,964	0.00775
60-64	558,224	0.0251	6,548	0.01173
65-69	400,320	0.0180	7,514	0.01877
70-74	302,464	0.0136	8,584	0.02838
75-79	215,728	0.0097	9,212	0.04270
80-84	117,872	0.0053	8,877	0.07531
85-89	53,376	0.0024	6,816	0.12769
90-94	15,568	0.0007	3,241	0.20820
95-99	2,224	0.0001	724	0.32576
100+	2,224	0.0001	1,089	0.48975
<b>Total</b>	<b>22,240,000</b>	<b>0.9951</b>	<b>102,449</b>	<b>1.37655</b>

CDR per 1,000

4.61




# Population distribution by country

Age group	United States (%)	Venezuela (%)	Ratio United States / Venezuela
0-1	1.39	2.19	0.63
1-4	5.48	8.00	0.69
5-9	6.75	10.38	0.65
10-14	7.19	10.23	0.70
15-19	7.24	10.12	0.72
20-24	7.02	9.25	0.76
25-29	6.71	8.42	0.80
30-34	6.79	7.26	0.94
35-39	7.06	6.88	1.03
40-44	7.66	6.30	1.22
45-49	7.57	5.15	1.47
50-54	6.75	4.37	1.54
55-59	5.75	3.46	1.66
60-64	4.37	2.51	1.74
65-69	3.39	1.80	1.88
70-74	2.84	1.36	2.09
75-79	2.48	0.97	2.56
80-84	1.87	0.53	3.53
85-89	1.07	0.24	4.46
90-94	0.45	0.07	6.43
95-99	0.13	0.01	13.00
100+	0.03	0.01	3.00
<b>Total</b>	<b>99.99</b>	<b>99.51</b>	<b>1.00</b>


- The U.S. has an older population than Venezuela
- This is causing  $CDR_{US} > CDR_{VE}$




# Age structure



# The U.S. has an older population than Venezuela



# Age-specific death rates, 2006



# Standardize Venezuela's *CDR*

Standardization Age group	Venezuela (observed rates)	United States (standard prop. population)	Venezuela's rates times U.S. prop. population
0-1	0.01623	0.0139	0.0002
1-4	0.00064	0.0548	0.0000
5-9	0.00032	0.0675	0.0000
10-14	0.00041	0.0719	0.0000
15-19	0.00146	0.0724	0.0001
20-24	0.00238	0.0702	0.0002
25-29	0.00221	0.0671	0.0001
30-34	0.00203	0.0679	0.0001
35-39	0.00215	0.0706	0.0002
40-44	0.00289	0.0766	0.0002
45-49	0.00411	0.0757	0.0003
50-54	0.00568	0.0675	0.0004
55-59	0.00775	0.0575	0.0004
60-64	0.01173	0.0437	0.0005
65-69	0.01877	0.0339	0.0006
70-74	0.02838	0.0284	0.0008
75-79	0.04270	0.0248	0.0011
80-84	0.07531	0.0187	0.0014
85-89	0.12769	0.0107	0.0014
90-94	0.20820	0.0045	0.0009
95-99	0.32576	0.0013	0.0004
100+	0.48975	0.0003	0.0001
<b>Total</b>		<b>0.9999</b>	<b>0.0097</b>

CDR per 1,000

9.68



# Another way... same results...

Standardization Age group	United States (standard population)	Venezuela (observed rates)	Venezuela (standardized deaths)
0-1	4,148,175	0.0162	67,325
1-4	16,353,955	0.0006	10,467
5-9	20,144,014	0.0003	6,446
10-14	21,457,106	0.0004	8,797
15-19	21,606,321	0.0015	31,545
20-24	20,949,775	0.0024	49,860
25-29	20,024,642	0.0022	44,254
30-34	20,263,386	0.0020	41,135
35-39	21,069,147	0.0022	45,299
40-44	22,859,726	0.0029	66,065
45-49	22,591,139	0.0041	92,850
50-54	20,144,014	0.0057	114,418
55-59	17,159,716	0.0078	132,988
60-64	13,041,384	0.0117	152,975
65-69	10,116,772	0.0188	189,892
70-74	8,475,408	0.0284	240,532
75-79	7,401,060	0.0427	316,025
80-84	5,580,638	0.0753	420,278
85-89	3,193,199	0.1277	407,740
90-94	1,342,934	0.2082	279,599
95-99	387,959	0.3258	126,381
100+	89,529	0.4898	43,847
<b>Total</b>	<b>298,400,000</b>		<b>2,888,718</b>

CDR per 1,000


9.68



# Comparing crude death rates

- $CDR_{\text{United States original}}$   
= 8.41 deaths per 1,000
- $CDR_{\text{Venezuela original}}$   
= 4.61 deaths per 1,000
- $CDR_{\text{Venezuela standardized}}$   
= 9.68 deaths per 1,000





**ATM** | **TEXAS A&M**  
UNIVERSITY.

# Life table

- One of the most important and elegant measures of the mortality experiences of a population is the life table
- It dates back to John Graunt (1620–1674) and his “Bills of Mortality”
- Demographers use the life table to determine life expectancy, not only at birth but at any age

# Important information from life table

- Like the total fertility rate (*TFR*), the life table is a synthetic or hypothetical measure
- It tells us how many years of life, on average, may a person expect to live if the person during his or her lifetime is subjected to the age-specific probabilities of dying of a particular country or population at a given time

# Example: United States, 2010

- Let's say that the population of the U.S. in 2010 had a life expectancy of birth of 78.7 years
- This means that if a cohort of persons, throughout the years of their life, were subjected to the *ASDRs* ( ${}_nM_x$ ) of the total population in the U.S. in 2010
  - They would live, on average, 78.7 years

# Examples of Life expectancy

- Life expectancy at birth is a primary indicator of quality of life

2013	Life expectancy	
	Male	Female
World	69	73
More developed countries	75	82
Less developed countries (except China)	65	69
Japan	80	86
Lesotho	42	45
Sierra Leone	45	46



# Limitations of $e_0$

- We need to be aware of the fact that when considering life expectancy at birth ( $e_0$ ), infant mortality plays a very important role
- When  $e_0$  is low, a major reason is their very high infant mortality rate
- When comparing values of life expectancy at birth across countries, we should not think of  $e_0$  as a modal age at death

# Abridged life table

- An abridged life table is calculated for age groups
- Usually for five-year age groups
- Rather than for single-year age groups

# Radix and mortality probabilities

- A life table starts with a population (a radix) of 100,000 persons born alive at age 0
  - This number is arbitrary, but conventional
  - It can also be 1,000 or 1
- From each age to the next, the population is decremented according to age-specific mortality probabilities until all members have died
  - The mortality schedule is fixed and does not change over the life of the population



# Life table for U.S. population, 2010

Life Table for the Total Population, United States, 2010						
(1) Age range	(2) ${}_nq_x$	(3) $l_x$	(4) ${}_nd_x$	(5) ${}_nL_x$	(6) $T_x$	(7) $e_x$
<1	0.006123	100,000	612	99,465	7,866,027	78.7
1-4	0.001071	99,388	106	397,294	7,766,561	78.1
5-9	0.000573	99,281	57	496,250	7,369,267	74.2
10-14	0.000708	99,224	70	495,989	6,873,017	69.3
15-19	0.002463	99,154	244	495,240	6,377,028	64.3
20-24	0.004317	98,910	427	493,529	5,881,789	59.5
25-29	0.004791	98,483	472	491,249	5,388,260	54.7
30-34	0.005497	98,011	539	488,744	4,897,011	50.0
35-39	0.006913	97,472	674	485,753	4,408,267	45.2
40-44	0.009979	96,798	966	481,758	3,922,514	40.5
45-49	0.016044	95,833	1,538	475,584	3,440,756	35.9
50-54	0.024343	94,295	2,295	466,066	2,965,173	31.4
55-59	0.035106	92,000	3,230	452,347	2,499,106	27.2
60-64	0.049847	88,770	4,425	433,348	2,046,759	23.1
65-69	0.074406	84,345	6,276	406,912	1,613,411	19.1
70-74	0.112315	78,069	8,768	369,612	1,206,499	15.5
75-79	0.174782	69,301	12,113	317,694	836,886	12.1
80-84	0.274384	57,188	15,692	248,038	519,193	9.1
85-89	0.430820	41,497	17,878	162,723	271,155	6.5
90-94	0.615282	23,619	14,532	79,720	108,432	4.6
95-99	0.783397	9,087	7,119	24,670	29,212	3.2
100+	1.00000	1,968	1,968	4,542	4,542	2.3

Source: Arias (2014: 62).



# Basic life table columns

1. Age intervals of each group
2.  ${}_nq_x$ : probability of dying between age  $x$  and age  $x+n$
3.  $l_x$ : number of survivors at each age  $x$
4.  ${}_nd_x$ : number of deaths between age  $x$  and age  $x+n$
5.  ${}_nL_x$ : number of years lived by all persons who enter the age interval while in the age interval
6.  $T_x$ : number of years lived by the population in the age interval and in all subsequent intervals
7.  $e_x$ : remaining life expectancy at each age



# 1. Age intervals of each group

- Age groups refer to the range of years between two birthdays
- The age group 5–9 refers to the five-year interval between the fifth and the tenth birthdays

## 2. Probabilities of dying ( ${}_nq_x$ )

- The most basic column of the life time shows probabilities of dying for each age group ( ${}_nq_x$ )
  - These are probabilities that persons alive at the beginning of an age interval will die during the interval, before they reach the start of the next age interval

$${}_nq_x = {}_nd_x / l_x$$

- For last age group,  ${}_nq_x=1.0$  because everybody dies

# Rates and probabilities

- Difference between mortality rates ( ${}_nM_x$ ) and mortality probabilities ( ${}_nq_x$ ) is the denominator
- ${}_nM_x$ : denominator is midyear population
- ${}_nq_x$ : denominator is population alive at the beginning of the age interval

### 3. Number of survivors ( $l_x$ )

- Number of people alive at the beginning of the age interval ( $l_x$ )
  - Known as “the little l column”
- It is calculated by subtracting the number of people dying ( ${}_nd_x$ ) from the  $l_x$  value in the age interval immediately preceding the one being calculated
- Example of U.S. life table in 2010
  - Of the 99,224 people alive at the beginning of the age interval 10–14 ( $l_{10}$ )
  - 70 of them die during the age interval ( ${}_5d_{10}$ )
  - Thus, the value of  $l_{15}$  is  $99,154 = 99,224 - 70$



## 4. Number of deaths ( ${}_nd_x$ )

- Number of people who die during a particular age interval ( ${}_nd_x$ )

$${}_nd_x = l_x * {}_nq_x$$

- For the number of people who die during the age interval of 40–44

$${}_5d_{40} = {}_5q_{40} * l_{40}$$

$${}_5d_{40} = 0.009979 * 96,798$$

$${}_5d_{40} = 966$$



## 5. Years lived in age interval ( ${}_nL_x$ )

- Total number of years lived by all persons who enter that age interval while in the age interval ( ${}_nL_x$ )
  - Known as “the big L column”
- Example of U.S. life table in 2010
  - 98,011 persons are alive at the beginning of age interval 30–34 ( $l_{30}$ )
  - If none of them died during the age interval, they would have lived 490,055 years (98,011 times 5)
  - But 539 of them died ( ${}_5d_{30}$ )



# Different formulas for ${}_nL_x$

- Demographers assume that deaths are roughly distributed during the five-year period for many of the age intervals
- This assumption does not apply to the first few age intervals
  - There are several formulas to produce the  ${}_nL_x$  value for the first few age groups
- At the other age extreme, 100+ in the life table, another formula is used



## 6. Years lived in current and subsequent age intervals ( $T_x$ )

- Total number of years lived by the population in the age interval and in all subsequent age intervals ( $T_x$ )
  - We sum  ${}_nL_x$  from the oldest age backwards to get  $T_x$

$$T_x = \sum_{i=x}^w L_i$$

- $L_i$ : entry  $i$  in the  ${}_nL_x$  column
- $\sum_{i=x}^w$  : sum of the  ${}_nL_x$  column starting at entry  $x$  through the last  ${}_nL_x$  entry ( $w$ )

# Example of $T_x$

- Example of U.S. life table in 2010

$$T_{95} = {}_5L_{95} + {}_{\infty}L_{100}$$

$$T_{95} = 24,670 + 4,542$$

$$T_{95} = 29,212$$



# 7. Remaining life expectancy ( $e_x$ )

- Average number of years of life remaining at the beginning of the age interval ( $e_x$ )
- It provides life expectancy at any age

$$e_x = T_x / l_x$$

- Example of U.S. life table in 2010

$$e_0 = T_0 / l_0 = 7,866,027 / 100,000 = 78.7$$

$$e_{25} = T_{25} / l_{25} = 5,388,260 / 98,483 = 54.7$$





- Persons aged 25–29 can expect to live an additional 54.7 years



# Average age at death ( $x + e_x$ )


- $e_x$  is the expectation of future life beyond age  $x$ 
  - It is not an average age at death
- We add  $x$  and  $e_x$  to obtain the average age at death for cohort members who survive to age  $x$ 
  - Not all lifetables include  $x + e_x$
  - The  $x + e_x$  column always goes up
- $e_x$  does not always go down
  - It often goes up after the first few years of life, because babies who survive infancy are no longer subject to the high risks of infancy

# Typical shapes of lifetable functions



# Stable population

- Alfred Lotka (1880–1949) used life tables in the development of his stable population theory
- If a population that is closed to migration experiences constant schedules of age-specific fertility and mortality rates
  - It will develop a constant age distribution
  - It will grow at a constant rate, irrespective of its initial age distribution



**ATM** | **TEXAS A&M**  
UNIVERSITY.

# Life table (extra)

- Estimate overall mortality of population
  - **Assumption**: age-specific rates for the period continue unchanged into the future
  - **Synthetic cohort**: imaginary cohort of new born babies would experience a life table from a specific period
  - **Life expectancy**: average age at death for a hypothetical cohort born in a particular year and being subjected to the risks of death experienced by people of all ages in that year

# Life table, U.S. women, 2010

Age interval	Number of females in the population	Number of deaths in the population	Age-specific death rates in the interval	Of 100,000 hypothetical people born alive:		Number of years lived		Expectation of life	
				Probabilities of death (proportion of persons alive at beginning who die during interval)	Number alive at beginning of interval	Number dying during age interval	In the age interval	In this and all subsequent age intervals	Average number of years of live remaining at beginning of age interval
$x$ to $x + n$	${}_nP_x$	${}_nD_x$	${}_nM_x$	${}_nq_x$	$l_x$	${}_nd_x$	${}_nL_x$	$T_x$	$e_x$
Under 1	1,976,387	11,503	0.00582	0.005791	100,000	579	99,508	8,098,622	81.0
1-4	7,905,548	1,976	0.00025	0.000999	99,421	99	397,445	7,999,114	80.5
5-9	9,959,019	1,095	0.00011	0.000550	99,322	55	496,471	7,601,670	76.5
10-14	10,097,332	1,313	0.00013	0.000650	99,267	65	496,173	7,105,199	71.6
15-19	10,736,677	3,436	0.00032	0.001599	99,202	159	495,615	6,609,025	66.6
20-24	10,571,823	4,757	0.00045	0.002247	99,044	223	494,662	6,113,410	61.7
25-29	10,466,258	5,652	0.00054	0.002696	98,821	266	493,440	5,618,747	56.9
30-34	9,965,599	6,876	0.00069	0.003444	98,555	339	491,925	5,125,308	52.0
35-39	10,137,620	10,138	0.00100	0.004988	98,215	490	489,852	4,633,382	47.2
40-44	10,496,987	17,005	0.00162	0.008067	97,725	788	486,656	4,143,531	42.4
45-49	11,499,506	29,094	0.00253	0.012570	96,937	1,219	481,639	3,656,874	37.7
90-94	1,023,979	165,495	0.16162	0.575549	29,621	17,048	105,484	148,164	5.0
95-99	288,981	78,398	0.27129	0.808265	12,573	10,162	37,458	42,680	3.4
100+	44,202	20,403	0.46159	1.000000	2,411	2,411	5,222	5,222	2.2



# Probability of dying ( ${}_nq_x$ )

- Need to convert age-specific death rates ( ${}_nM_x$ ) to probabilities of dying ( ${}_nq_x$ )
- Probability of death relates the number of deaths during any given number of years to the number of people who started out being alive and at risk of dying

$${}_nq_x = (n)({}_nM_x) / 1 + (a)(n)({}_nM_x)$$

- $(a)(n)$ : average years lived per person by people dying in the interval.  $a=0.5$  implies that deaths are distributed evenly over an age interval. For 0–1 age,  $a=0.85$ . For 1–4 age,  $a=0.60$ .
- For last group,  $q=1.0$ .

# Number of deaths ( ${}_nd_x$ ) and alive ( $l_x$ )

- The life table assumes an initial population of 100,000 births (radix), which is subjected to the mortality schedule
  - Radix can also be 1
- Number of people dying during age interval ( ${}_nd_x$ ) equals probability of death times number alive at beginning ( $l_x$ )

$${}_nd_x = ({}_nq_x)(l_x)$$

- Subtracting those who died in the previous age interval gives the number of people still alive at the beginning of next age interval

$$l_{x+n} = l_x - {}_nd_x$$



# Number of years lived ( ${}_nL_x$ )

- Number of years lived ( ${}_nL_x$ ) has to consider that some people die before the end of the age interval
- The lower the death rates, more people will survive through an entire age interval

$${}_nL_x = n(l_x - a_nd_x)$$

- $a$ : usually 0.5, which implies that deaths are distributed evenly over an age interval. For 0–1 age,  $a=0.85$ . For 1–4 age,  $a=0.60$
- ${}_nL_x$  for the oldest, open-age interval

$$L_{100+} = l_{100} / M_{100}$$

- $l_{100}$ : number of survivors to oldest age
- $M_{100}$ : death rate at the oldest age



# Cumulative number of years lived ( $T_x$ )

- Number of years lived are added up, cumulating from the oldest to the youngest ages
- Total number of years lived in a given age interval and all older age intervals ( $T_x$ )

$$T_x = T_{x+n} + {}_nL_x$$

- At the oldest age,  $T_x$  equals  ${}_nL_x$

# Life expectancy ( $e_x$ )

- Expectation of life is the average remaining lifetime
- It is the total years remaining to be lived at exact age  $x$
- Division of total number of years lived ( $T_x$ ) by number of people alive at that exact age ( $I_x$ )

$$e_x = T_x / I_x$$

- This index summarizes the level of mortality prevailing in a given population at a particular time

# Probability of surviving ( $p_x$ )

- Probability of surviving from birth to age  $x$  is designated  $p_x$

$$p_x = l_x / l_0$$

- We can also estimate the probability of surviving from one particular age group to the subsequent age group

# Crude death and birth rates

- Crude death rate (CDR) equals total number of deaths ( $I_0$ ) divided by total population ( $T_0$ )
- Crude birth rate (CBR) equals total number of births ( $I_0$ ) divided by total population ( $T_0$ )

$$\text{CDR} = \text{CBR} = I_0 / T_0 = 1 / (T_0 / I_0) = 1 / e_0$$



# Alternative interpretations

- **Synthetic cohort** (history of a hypothetical cohort)
  - Lifetime mortality experience of a single cohort of newborn babies, who are subject to specific age-specific mortality rates
  - Used in public health/mortality studies, calculation of survival rates for estimating population, fertility, net migration...
- **Stationary population**
  - Results from unchanging schedule of age-specific mortality rates and a constant annual number of births/deaths (radix)
  - Used in the comparative measurement of mortality and in studies of population structure

# Same interpretation

- **$x$  to  $x+n$** 
  - Period of life between two exact ages
  - For instance, 20–25 means the 5-year interval between the 20<sup>th</sup> and 25<sup>th</sup> birthdays
- **${}_nq_x$** 
  - Proportion of persons in the cohort alive at the beginning of an indicated age interval ( $x$ ) who will die before reaching the end of that age interval ( $x+n$ )
  - Probability that a person at his/her  $x^{\text{th}}$  birthday will die before reaching his/her  $x+n^{\text{th}}$  birthday
- **$e_x$  (life expectancy)**
  - Average remaining lifetime (in years) for a person who survives to the beginning of the indicated age interval

$l_x$

- **Synthetic cohort**

- Number of persons living at the beginning of the indicated age interval (x) out of the total number of births assumed as the radix of the table

- **Stationary population**

- Number of persons who reach the beginning of the age interval each year

$${}_n d_x$$

- **Synthetic cohort**
  - Number of persons who would die within the indicated age interval (x to x+n) out of the total number of births assumed in the table
- **Stationary population**
  - Number of persons that die each year within the indicated age interval

$${}_nL_x$$

- **Synthetic cohort**

- Number of person-years that would be lived within the indicated age interval (x to x+n) by the cohort of 100,000 births assumed

- **Stationary population**

- Number of persons in the population who at any moment are living within the indicated age interval

$$T_x$$

- **Synthetic cohort**

- Total number of person-years that would be lived after the beginning of the indicated age interval by the cohort of 100,000 births assumed

- **Stationary population**

- Number of persons in the population who at any moment are living within the indicated age interval and all higher age intervals



# Interpretation as stationary population

ABRIDGED LIFE TABLE FOR THE FEMALE POPULATION OF THE UNITED STATES: 2007

Age group	Width n	Population nPx	Deaths nDx	Of 100,000 born alive				Stationary population		Average remaining lifetime ex
				Age-specific death rates nMx	Proportion dying nqx	# living at beginning of interval lx	# dying during interval ndx	In the age interval nLx	In this and following ages Tx	
0	1	1,998,761	12,845	0.0064	0.0064	100,000	641	99,684	8,103,588	81.0
1-4	4	8,109,371	2,069	0.0003	0.0010	99,359	101	397,248	8,003,904	80.6
5-9	5	9,720,587	1,192	0.0001	0.0006	99,258	61	496,150	7,606,656	76.6
10-14	5	9,918,543	1,370	0.0001	0.0007	99,197	68	495,828	7,110,506	71.7
15-19	5	10,617,178	3,741	0.0004	0.0018	99,129	175	495,242	6,614,678	66.7
20-24	5	10,073,754	4,925	0.0005	0.0024	98,954	242	494,215	6,119,436	61.8
25-29	5	10,122,681	5,824	0.0006	0.0029	98,713	284	492,910	5,625,222	57.0
30-34	5	9,469,789	6,956	0.0007	0.0037	98,429	361	491,314	5,132,312	52.1
35-39	5	10,666,827	11,126	0.0010	0.0052	98,068	510	489,165	4,640,998	47.3
40-44	5	11,155,652	18,375	0.0016	0.0082	97,558	800	485,944	4,151,834	42.6
45-49	5	11,572,428	29,834	0.0026	0.0128	96,757	1,240	480,926	3,665,890	37.9
50-54	5	10,709,011	40,396	0.0038	0.0187	95,518	1,786	473,463	3,184,963	33.3
55-59	5	9,339,919	50,868	0.0054	0.0269	93,732	2,521	462,827	2,711,501	28.9
60-64	5	7,636,068	62,624	0.0082	0.0402	91,211	3,670	447,543	2,248,674	24.7
65-69	5	5,725,079	74,499	0.0130	0.0631	87,541	5,528	424,827	1,801,131	20.6
70-74	5	4,738,379	96,395	0.0203	0.0971	82,012	7,962	391,395	1,376,304	16.8
75-79	5	4,314,403	139,360	0.0323	0.1500	74,050	11,109	343,929	984,910	13.3
80-84	5	3,582,388	192,519	0.0537	0.2378	62,941	14,970	278,566	640,981	10.2
85+	---	3,511,395	464,781	0.1324	1.0000	47,971	47,971	362,415	362,415	7.6


ABRIDGED LIFE TABLE FOR THE MALE POPULATION OF THE UNITED STATES: 2007

Age group	Width n	Population nPx	Deaths nDx	Of 100,000 born alive				Stationary population		Average remaining lifetime ex
				Age-specific death rates nMx	Proportion dying nqx	# living at beginning of interval lx	# dying during interval ndx	In the age interval nLx	In this and following ages Tx	
0	1	2,079,846	16,293	0.0078	0.0078	100,000	780	99,615	7,582,342	75.8
1-4	4	8,507,893	2,634	0.0003	0.0012	99,220	123	396,648	7,482,726	75.4
5-9	5	10,095,353	1,519	0.0002	0.0008	99,097	75	495,313	7,086,078	71.5
10-14	5	10,484,813	2,066	0.0002	0.0010	99,022	98	494,887	6,590,765	66.6
15-19	5	11,252,863	9,558	0.0008	0.0042	98,925	419	493,658	6,095,878	61.6
20-24	5	10,828,130	15,758	0.0015	0.0073	98,505	714	490,881	5,602,220	56.9
25-29	5	10,489,470	15,107	0.0014	0.0072	97,791	702	487,338	5,111,340	52.3
30-34	5	9,802,132	14,685	0.0015	0.0075	97,089	725	483,776	4,624,002	47.6
35-39	5	10,684,227	19,755	0.0018	0.0092	96,364	887	479,777	4,140,226	43.0
40-44	5	11,085,591	30,350	0.0027	0.0136	95,477	1,299	474,390	3,660,450	38.3
45-49	5	11,318,167	47,904	0.0042	0.0210	94,179	1,974	466,332	3,186,060	33.8
50-54	5	10,313,298	66,552	0.0065	0.0318	92,205	2,931	454,237	2,719,728	29.5
55-59	5	8,790,943	81,590	0.0093	0.0454	89,274	4,055	436,954	2,265,491	25.4
60-64	5	6,979,426	92,028	0.0132	0.0640	85,218	5,451	413,393	1,828,537	21.5
65-69	5	5,003,042	100,492	0.0201	0.0959	79,767	7,651	380,904	1,415,144	17.7
70-74	5	3,889,104	117,852	0.0303	0.1414	72,116	10,196	336,467	1,034,240	14.3
75-79	5	3,192,676	149,669	0.0469	0.2107	61,920	13,046	278,295	697,773	11.3
80-84	5	2,235,826	171,134	0.0765	0.3220	48,874	15,739	205,629	419,478	8.6
85+	---	1,606,146	248,866	0.1549	1.0000	33,135	33,135	213,850	213,850	6.5


Source: Formulas from Kintner (2003); Population data from 2007 ACS; Death data from CDC ([http://www.cdc.gov/nchs/data/dvs/mortfinal2007\\_worktable310.pdf](http://www.cdc.gov/nchs/data/dvs/mortfinal2007_worktable310.pdf)).



# Population, U.S., 2007



# ${}_nL_x$ from previous life tables, U.S., 2007




# Problems with life tables

- We saw life tables based on complete empirical data
- We might experience some issues
  - Have partial information to build our life table
  - Have data for only some age groups
  - Information for some ages may be more reliable than for other ages
  - Have ideas about mortality level, but not a full life table to make projections
- We can use model life tables to solve these issues

# Model life tables

- A life table constructed from mathematical formulas is called a model life table
  - Use mathematical formulas to fill in missing parts
  - Have a whole life table from partial information
  - Identify suspicious and poor quality data with model expectations
  - Supply standard assumptions for projections
  - Find regularities for the invention of indirect measures
  - Reconstruct rates from historical counts of births and deaths (inverse projection)



**ATM** | **TEXAS A&M**  
UNIVERSITY.

# Short history of mortality in the world

- Defining the health and mortality transition
- Health and mortality changes over time
- Life span and longevity
- Disease and death over the life cycle
- Causes of poor health and death
- Health and mortality inequalities



# Morbidity and mortality

- Health, death: two sides of morbidity, mortality
  - Morbidity: prevalence of disease in a population
  - Mortality: pattern of death
- Health and mortality transition
  - Epidemiologic transition (Omran 1971)
  - Shift from prevailing poor health (high morbidity) and high death rates (high mortality) primarily from communicable diseases, occurring especially among the young...
  - To prevailing good health and low deaths rates from infectious diseases, with most people dying at older ages from degenerative diseases




# Death at older ages

- For virtually all of human history, early death was commonplace
- Beginning about 200 years ago, we have been steadily pushing death to older ages
- The survival of more people to ever older ages is a key contribution to the demographic transition
- Most people now survive to advanced ages and die pretty quickly
- The variability by age in mortality is compressed, leading to an increased rectangularization of mortality...



# Rectangularization of mortality, United States




# Changes over time

- For most of history, life expectancy fluctuated between 20 and 30 years
- About 2/3 of babies survived to their first birthday, and about 1/2 were still alive at age five
  - Now it's 99%
- Around 10% of people made it to age 65 in a pre-modern society
  - Now it's 90%



# Female life expectancy at birth



# Health improvements

Period and regions	Life expectancy (females)	% surviving to age				Births required for ZPG
		1	5	25	65	
Premodern	20	63	47	34	8	6.1
	30	74	61	50	17	4.2
US/Europe, late 18 <sup>th</sup> /early 19 <sup>th</sup>	40	82	73	63	29	3.3
Lowest Sub-Saharan	46	89	82	75	34	2.7
World average circa 2015	73	98	98	97	77	2.1
Mexico	78	99	99	98	84	2.1
United States	81	99	99	99	88	2.1
Canada	84	99	99	99	91	2.1
Japan (highest in the world)	86	99	99	99	93	2.1



# The Roman era

- Life expectancy in the Roman era is estimated to have been 22 years
- People who reached adulthood were not too likely to reach a very advanced age, although of course some did



# The Middle Ages

- The plague (black death) hit Europe in the 14<sup>th</sup> century, having spread west from Asia
  - An estimated 1/3 of the population of Europe may have perished from the disease between 1346 and 1350
- It appears to be the same disease that exists today
  - Not really known why it was so fatal back then
  - Probably due to generally poor health and few resources to battle the disease



# The Columbian exchange

- Columbus and other European explorers took diseases, horses, and guns to the Americas
  - Brought back new foods and few new diseases
- One explanation for relative ease with which Spain dominated Latin America after arriving around 1500
  - Explorers had immunity to the diseases they brought
  - Compared to the devastation the diseases affected indigenous populations



# Industrial Revolution, 1760–1840

- Plague and Little Ice Age had receded
- Income improved nutrition, housing, and sanitation
- Life expectancy in Europe and the U.S.:  $\approx 40$  years
  - Was population growth a cause or effect of rising living standards?
- There were as many deaths to children under 5 as there were at 65 and over
  - Infectious diseases were still the dominant reasons for death, but their ability to kill was diminishing



# 19<sup>th</sup> century

- Key elements in postponing death
  - Belief in the power of human intervention (Western science)
  - Improved nutrition: occurred first in Western Europe
  - Clean water, toilets, bathing facilities
  - Sewerage in cities: sanitation studies in Liverpool
  - Small pox vaccinations: Edward Jenner in England
- Validation of germ theory
  - Ignaz Semmelweis in Vienna: pioneer of antiseptic procedures
  - Joseph Lister in Glasgow: cleanliness principals in surgery
  - Louis Pasteur in Paris: formal experiments about germs, diseases




# 20<sup>th</sup> century

- Health as a social movement
  - Leading to government-organized universal health care systems in all rich countries except the U.S.
- Antibiotics emerging around WWII
- More vaccinations
- Oral rehydration therapy for infants and adults
- Advanced diagnoses, drugs, and other treatments for degenerative diseases to keep older people alive longer




# Improvements in life expectancy



	1850	1860	1870	1880	1890	1900	1910	1920	1930	1940	1950	1960	1970	1980	1990	2000	2010
Mexico						29.8	25.4	31.5	36.7	45.7	53.7	59.5	62.7	70.3	74.5	76.9	77.7
United States	40.5	40.9	42	43.5	44.5	48.3	51.8	54.6	61.6	65.2	71.1	73.1	74.8	77.4	78.8	79.6	80.9
Canada								60.6	62.1	66.3	70.9	74.3	76.5	79.1	80.4	81.9	83.5



# World War II: a turning point



# Postponing death

- Two ways to postpone death to the oldest possible ages
- Prevent diseases from occurring or spreading when they do occur (prevention)
  - Vaccinations, clean water, sanitation, good nutrition
  - No physicians needed
- Curing people of disease when they are sick
  - Diagnostic technology, drugs, skilled physicians



# Nutrition transition and obesity

- Poor were skinny because only the rich could afford to access enough quantities of food
  - Not any more
- Nutrition transition is a worldwide shift toward
  - Diet high in fat and processed foods
  - Diet low in fiber
  - Less exercise
  - Increases in degenerative diseases



# Life span and longevity

- Life span: oldest age to which human beings can survive
  - 122 years and 164 days (by 2016)
  - It is almost entirely a biological phenomenon
- Longevity/life expectancy: age at which we actually die
  - Expected number of years to be lived, on average, by a particular population at a particular time
  - Currently about 71 for all humans
  - It has biological and social components
- Populations with high mortality tend to have high morbidity
  - This is not a one-to-one relationship
  - We may live longer even though not being very healthy




# Mortality by age

- Humans are like most other animals with respect to the general pattern of death by age
  - The very young and the old are most vulnerable
  - Young adults are least likely to die
- Risks of death are relatively low
  - After the initial year of life, lasting at least until middle age
  - Corresponds to reproductive ages
- Beyond middle age
  - Mortality increases
  - Although at a decelerating rate



# Highest death rates, 2011: very young and the old



# Mortality by sex (gender)


- Women (sex)
  - Have a lower probability of death at every age from the moment of conception'
- Women (gender)
  - Unless society intervenes with a lower status for women that gives them less food, less access to health care, less education, lower earnings...



# Other mortality differentials

- Urban and rural differentials
  - Urban now better than rural
- Neighborhood inequalities
  - Slums are bad for your health
- Educational differentials
  - Better educated live longer
- Social status differentials
  - The rich live longer
- Race and ethnicity differentials
  - Being different will be used against you
- Marital status
  - Being married is good for your health





**ATM** | **TEXAS A&M**  
UNIVERSITY.

# Mortality trends and causes of death

- Categories of diseases according to the World Health Organization
  - Communicable diseases
  - Noncommunicable diseases
  - Injuries


# Communicable diseases

- Bacterial (e.g., tuberculosis)
- Viral (e.g., measles)
- Protozoan (e.g., malaria)
- Maternal conditions
  - Lack of prenatal care
  - Delivering somewhere besides a hospital
  - Seeking an unsafe abortion
- Perinatal conditions
  - “Surrounding birth” — just before and just after birth
- Nutritional deficiencies



Cause of death	Broad category of cause	Number of deaths in world 2011 (millions)	Top ten death rates (per 100,000 population), 2011			
			High income countries	Upper middle income countries	Lower middle income countries	Low income countries
Ischemic heart disease	Non-Com.	7.0	119	120	93	47
Stroke	Non-Com.	6.2	69	126	75	56
Lower respiratory infection	Com.	3.2	32	22	60	98
Chronic obstructive pulmonary disease (COPD)	Non-Com.	3.0	32	45	51	
Diarheal diseases	Com.	1.9			47	69
HIV/AIDS	Com.	1.6			24	70
Trachea bronchus, lung cancers	Non-Com.	1.5	51	28		
Diabetes mellitus	Non-Com.	1.4	21	20	20	
Road injury	Injury	1.3		21	19	
Prematurity	Com.	1.2			27	43
Alzheimer's disease and other dementias	Non-Com.		48			
Colon rectal cancers	Non-Com.		27			
Hypertensive heart disease	Non-Com.		20	18		
Breast cancer	Non-Com.		16			
Malaria	Com.					38
Tuberculosis	Com.				22	32
Protein-energy malnutrition	Com.					32
Birth asphyxia and birth trauma	Com.					30
Liver cancer	Non-Com.			19		
Stomach cancer	Non-Com.			18		
Life expectancy at birth (both sexes)			80	74	66	60

## The 10 leading causes of death in the world 2012



Source: World Health Organization. Available at:  
<http://www.who.int/mediacentre/factsheets/fs310/en/> (accessed April 29, 2016)

# Adults and children estimated to be living with HIV, 2013




Total: 35.0 million


95% confidence interval: [33.2 million – 37.2 million]

# Top ten causes of death, 2008

## Low-income countries



## Middle-income countries



## High-income countries






Figure 7.6. Life Expectancy at Birth: World and More developed, Less developed, and Least developed regions, 1950 to 2100



Source: United Nations, 2013d: 16. |



TEXAS A&M  
UNIVERSITY.

# Mortality and longevity in the U.S.

- Mortality started dropping gradually
  - In response to changes in the socioeconomic conditions and the environment of modernization
  - Much of the mortality reduction started to happen before the initiations of any appreciable public health measures
- Life expectancy increased
  - 46 for males and 48 for females in 1900
  - 76 for males and 81 for females in 2013



# Mortality in the United States


- Most improvements happened from 1900 to 1950
  - Germ theory: control of infectious and parasitic diseases
  - Boiling bottles and milk, washing hands, protecting food from flies, isolating sick children, ventilating rooms, improving water supply, sewage disposal
- Since 1950s, life expectancy improvements is due to prevention and control of chronic diseases
  - Heart disease, stroke...



# Hispanic paradox

- The Hispanic epidemiological paradox is the empirical finding that Hispanics have death rates of about the same magnitude as, and sometimes lower than, whites
- Also known as the Latino mortality paradox
- These findings are more evident for those of Mexican origin

# Life expectancy at birth by race/ethnicity: U.S., 2006–2011



# Explaining the Hispanic paradox

- Data artifacts
  - **Underreporting** of Hispanic-origin identification on death certificates
  - **Misstatement of age**, perhaps overstatement, at the older ages
- Migration effects
  - **Healthy migrant effect**: self-selection of immigrants in better physical and mental health
  - **Salmon bias**: Mexican Americans in poor health return to Mexico at old ages (return migrant effect)
- Cultural effects
  - **Better dietary** practices of Latinos and **stronger family** obligations and relationships



# Racial crossover

- Life expectancy at birth is the lowest for blacks compared with Hispanics and whites
- For most of their lives, blacks have higher death rates than Hispanics and whites
- The situation changes at the very oldest ages
- By late life, death rates for blacks become lower than those for whites, and in some cases lower than those for Hispanics




# Life expectancy at ages 70, 80, 90, and 100 by race/ethnicity and sex: United States, 2010


Age	Hispanics		NH-Whites		NH-Blacks	
	Males	Females	Males	Females	Males	Females
70	15.4	18.0	14.2	16.4	12.8	15.7
80	9.0	10.8	8.1	9.6	7.8	9.6
90	4.5	5.4	4.0	4.8	4.4	5.2
100	2.3	2.6	2.0	2.3	2.5	2.8

# Explaining the racial crossover

- Age misreporting on death certificates
  - Overstatement of age
  - But this would only postpone crossover to later ages, not eliminating it
- Population heterogeneity in frailty
  - The surviving elderly black population is a more robust group of disadvantaged individuals
  - The more frail blacks die before the age of 80 or 90
  - This produces a more robust group of blacks that live longer than the majority

# Percent distribution of five leading causes of death by age group: United States, 2011






TEXAS A&M  
UNIVERSITY.

# Infant mortality

- The infant mortality rate (IMR) is the most common measure of infant death
  - It is the number of deaths in a year to children under age 1 per 1,000 babies born in the year
- Declining infant mortality is key to population growth
- Reduction attributable especially to the development of oral rehydration therapy (ORT)
  - A solution of salts and sugars taken orally
  - Treats diarrhea: A major cause of death in young children
  - Developed in labs, tested in the field, especially Bangladesh
  - One of its founders still holds a teaching position at Harvard School of Public Health (Dr. Richard Cash)



# Infant mortality, 2015




**Table 7.4.** Countries with the Highest and the Lowest Infant Mortality Rates in the World, 2013

Highest infant mortality rates		Lowest infant mortality rates	
Central African Republic	116	Iceland	1.8
Congo, Dem. Rep.	109	Finland	1.8
Chad	96	Japan	1.9
Angola	96	Singapore	2.0
Guinea-Bissau	94	Estonia	2.1
Sierra Leone	92	Sweden	2.3

*Source:* Population Reference Bureau, 2014.

# Infant mortality rates by mother's race/ethnicity: United States, 2000–2010



# Neonatal and postneonatal mortality rates

- The infant mortality rate (IMR) may be thought of as the sum of two rates
- Neonatal mortality rate (NMR)
  - Deaths of babies aged 28 days or less per 1,000 live births
- Postneonatal mortality rate (PMR)
  - Deaths of babies aged between 29 days and 1 year per 1,000 live births



# Neonatal mortality rates and number of neonatal deaths

Region	Neonatal mortality rate (deaths per 1,000 live births)			Number of neonatal deaths (thousands)	
	1990	2013	Decline (percent) 1990–2013	1990	2013
<b>Developed regions</b>	8	3	55	118	48
<b>Developing regions</b>	36	22	40	4,554	2,714
<b>Northern Africa</b>	30	13	56	109	53
<b>Sub-Saharan Africa</b>	46	31	32	977	1,066
<b>Latin America and the Caribbean</b>	22	9	58	255	101
<b>Caucasus and Central Asia</b>	26	15	42	51	26
<b>Eastern Asia</b>	25	8	69	784	150
Excluding China	12	8	35	11	7
<b>Southern Asia</b>	51	30	42	1,940	1,086
Excluding India	49	30	39	578	338
<b>South-eastern Asia</b>	27	14	47	321	160
<b>Western Asia</b>	28	14	50	111	67
<b>Oceania</b>	26	21	19	5	6
<b>World</b>	<b>33</b>	<b>20</b>	<b>40</b>	<b>4,672</b>	<b>2,763</b>

# Endogenous and exogenous

- **Endogenous cause of death** in an infant can occur because of genetic issues or conditions associated with fetal development or the birth process
- **Exogenous cause of death** is due mainly to environmental or external factors, such as infections or accidents



# Causes of neonatal deaths

- The main causes of neonatal deaths are **endogenous conditions**
- Congenital malformations, chromosomal abnormalities, complications of delivery, low birthweight, genetic disorders...
- However, endogenous causes dominate infant mortality mainly in the early days of life, and not for the entire first month of life



# Causes of postneonatal deaths

- Postneonatal mortality rate was 18 per 1,000 live births for the world in 2013
  - Rate was 2 in developed countries
- Deaths in postneonatal period and in first few years of life are mainly due to **exogenous causes**
  - Infectious disease, accidents, injury
- Improved living standards, better healthcare, and public health programs have greater effects on exogenous causes than on endogenous causes



# Stillbirth rate (SBR)

- Stillbirths (miscarriages, fetal deaths)
  - A stillbirth is a fetus not born alive and is not registered as a death
  - SBR: stillbirths per 1,000 live births plus stillbirths in the year
  - Stillbirths are often identified in hospital reports dealing with obstetric procedures
- WHO: interventions can be planned if we know at what point before birth the fetus died
  - 2.6 million stillbirths in the world in 2009
  - 18.9 stillbirths per 1,000 live births plus stillbirths



# Perinatal mortality rate (PeMR)

- PeMR relates to stillbirths and deaths of babies who lived for only seven days or less per 1,000 live births plus stillbirths in the year
- **Endogenous** causes of mortality in the 1st week after birth are similar to the causes of stillbirths
- PeMR in 2010
  - World (47)
  - Developed world (10); less developed world (50)
  - Czech Republic and Singapore (4); Mauritania (111)
- U.S.: 6.5 in 2006; 6.3 in 2011



# Maternal mortality ratio (MMR)

- MMR measures the extent to which mothers die immediately before, during, or after giving birth because of problems associated with the pregnancy or childbirth
  - **Numerator:** number of deaths in a year of women dying as a result of complications of pregnancy, childbirth, and the puerperium (condition of the woman immediately following childbirth)
  - **Denominator:** live births occurring in the year
  - Multiplied by 100,000, because it is increasingly rare in developed countries



# Examples of maternal deaths


- World
  - 529,000 maternal deaths in 2000
  - 313,000 maternal deaths in 2015
- Developing regions in 2015
  - 99% of all maternal deaths
  - 66% in sub-Saharan Africa
  - 21% in southern Asia
- MMR in 2015
  - World: 216 per 100,000 live births
  - Sierra Leone: 1,360
  - Sub-Saharan Africa: 546
  - U.S., Iran, Hungary: 21
  - Greece, Singapore (3); Estonia (2)



# Factors associated with MMR

- Maternal deaths are mostly due to age, parity (number of children), birth spacing
  - Younger and older women are more likely, compared to women in their 20s and 30s
  - High-parity women are at high risk
  - Women with short birth intervals are also at high risk
- Other factors
  - Chronic disease and malnutrition, poverty, unwanted pregnancies, inadequate prenatal and obstetric care, lack of access to a hospital





TEXAS A&M  
UNIVERSITY.

# Future course of mortality

- Is it likely that mortality rates will continue to fall resulting in even higher levels of life expectancy than those attained by developed countries?
- There are two positions
  - One argues for a limit
  - The other argues against one
- Poston argues against a fixed biological maximum life expectancy (more optimistic view)



# Mortality reversals

- Modern medical and public health techniques will surely bring about further reductions in mortality from the major causes
- However, there is evidence since the early 1980s of mortality reversals
  - First in Eastern Europe
  - Later in sub-Saharan Africa

# Eastern Europe

- What is causing mortality reversals in Eastern Europe?
- Possible factors include
  - Lack of preventative health programs
  - Inadequate quality of medical services
  - Smoking and alcohol abuse
  - General neglect of individual health
- These were caused by
  - Lack of life choices under the former communist regimes
  - Unemployment, relative deprivation, and inability to cope with economic challenges of the capitalist regime



# Sub-Saharan Africa

- The AIDS epidemic has halted or reversed gains in life expectancy in many of the sub-Saharan African countries

# Degenerative diseases


- Degenerative diseases are the major causes of death in the developed world
- Future improvements in the treatment of these diseases is likely
  - However, eliminating a specific degenerative disease would result in small increases to life expectancy because these diseases occur to older individuals
  - Eliminating heart disease would raise life expectancy in the U.S. by 5 years, and eliminating cancer would raise it by 3 years



# Important question

- The important question is not “will life expectancy increase?”
- The important question is “by how much will life expectancy increase?”





**ATM** | **TEXAS A&M**  
UNIVERSITY.

# Coronavirus pandemic

- **Coronavirus COVID-19 Global Cases**

Center for Systems Science and Engineering (CSSE) at Johns Hopkins University (JHU)

<https://coronavirus.jhu.edu/map.html>

- **COVID-19 Coronavirus Pandemic**

Worldometer

<https://www.worldometers.info/coronavirus/>

- **COVID Community Vulnerability Map**

Jvion – Prescriptive Analytics for Preventable Harm

<https://covid19.jvion.com/>

- **Coronavirus Deaths by U.S. State and Country Over Time: Daily Tracking**

Josh Katz, Margot Sanger-Katz

<https://www.nytimes.com/interactive/2020/03/21/upshot/coronavirus-deaths-by-country.html?action=click&auth=login-email&login=email&module=Top%20Stories&pgtype=Homepage>



# Coronavirus and demography

- **Resources about demography and COVID-19**

Population Europe

<https://population-europe.eu/news/demography-coronavirus>

- **A demographer's view of the coronavirus pandemic**

Isaac Chotiner

March 13, 2020

<https://www.newyorker.com/news/q-and-a/a-demographers-view-of-the-coronavirus-pandemic>

- **Coronavirus: Why it's so deadly in Italy**

Andreas Backhaus

March 13, 2020

<https://medium.com/@andreasbackhausab/coronavirus-why-its-so-deadly-in-italy-c4200a15a7bf>

- **Demographic science aids in understanding the spread and fatality rates of COVID-19**

Jennifer Beam Dowd, Valentina Rotondi, Liliana Andriano, David M. Brazel, Per Block, Xuejie Ding, Yan Liu, Melinda C. Mills

March 14, 2020

[https://osf.io/fd4rh/?view\\_only=c2f00dfe3677493faa421fc2ea38e295&fbclid=IwAR15NPHaq\\_Ha9yK3uDDN0FhdAZmnfMLfdqgrHaL7qllf3jBYu8CA\\_9gcEq](https://osf.io/fd4rh/?view_only=c2f00dfe3677493faa421fc2ea38e295&fbclid=IwAR15NPHaq_Ha9yK3uDDN0FhdAZmnfMLfdqgrHaL7qllf3jBYu8CA_9gcEq)

- **Students could be undercounted in the census as coronavirus closes colleges – here's why that matters**

Dudley L. Poston, Jr.

March 23, 2020

<https://theconversation.com/students-could-be-undercounted-in-the-census-as-coronavirus-closes-colleges-heres-why-that-matters-133889>



# Coronavirus and sex

- **Sex, gender and COVID-19**

Global Health 5050

<http://globalhealth5050.org/covid19/>

- **Sex, gender and COVID-19: Disaggregated data and health disparities**

BMJ Global Health Blogs

March 24, 2020

<https://blogs.bmj.com/bmjgh/2020/03/24/sex-gender-and-covid-19-disaggregated-data-and-health-disparities/>

- **Here's why the coronavirus may be killing more men than women. The US should take note**

Katie Polglase, Gianluca Mezzofiore, Max Foster

March 24, 2020

<https://www.cnn.com/2020/03/24/health/coronavirus-gender-mortality-intl/index.html>



# Coronavirus and politics

- **The Real Pandemic Danger Is Social Collapse**

Branko Milanovic

March 19, 2020

[https://www.foreignaffairs.com/articles/2020-03-19/real-pandemic-danger-social-collapse?utm\\_medium=social&utm\\_source=twitter\\_cta&utm\\_campaign=cta\\_share\\_buttons](https://www.foreignaffairs.com/articles/2020-03-19/real-pandemic-danger-social-collapse?utm_medium=social&utm_source=twitter_cta&utm_campaign=cta_share_buttons)

- **Coronavirus Could Overwhelm U.S. Without Urgent Action, Estimates Say**

James Glanz, Lauren Leatherby, Matthew Bloch, Mitch Smith, Larry Buchanan, Jin Wu, Nicholas Bogel-Burroughs

March 20, 2020

<https://www.nytimes.com/interactive/2020/03/20/us/coronavirus-model-us-outbreak.html?action=click&module=Spotlight&pgtype=Homepage>

- **Before Virus Outbreak, a Cascade of Warnings Went Unheeded**

David E. Sanger, Eric Lipton, Eileen Sullivan, Michael Crowley

March 22, 2020

<https://www.nytimes.com/2020/03/19/us/politics/trump-coronavirus-outbreak.html>

- **Sisi and Erdogan Are Accomplices of the Coronavirus**

Steven A. Cook

March 17, 2020

<https://foreignpolicy.com/2020/03/17/sisi-and-erdogan-are-partners-with-the-coronavirus/>

- **How South Korea Flattened the Curve**

Max Fisher, Choe Sang-Hun

March 23, 2020


<https://www.nytimes.com/2020/03/23/world/asia/coronavirus-south-korea-flatten-curve.html?action=click&module=Top%20Stories&pgtype=Homepage>



# References

- Hviid A, Hansen JV, Frisch M, Melbye M. 2019. “Measles, mumps, rubella vaccination and autism: A nationwide cohort study.” *Annals of Internal Medicine*, March 5. (<http://www.doi.org/10.7326/M18-2101>)
- Poston DL, Bouvier LF. 2017. *Population and Society: An Introduction to Demography*. New York: Cambridge University Press. 2nd edition. Chapter 7 (pp. 163–214).
- Weeks JR. 2015. *Population: An Introduction to Concepts and Issues*. 12th edition. Boston: Cengage Learning. Chapter 5 (pp. 139–188).





TEXAS A&M  
UNIVERSITY.