

Lecture 15: Ordinary least squares regression

Ernesto F. L. Amaral

November 22–December 06, 2022
Advanced Methods of Social Research (SOCI 420)

Source: Healey, Joseph F. 2015. "Statistics: A Tool for Social Research." Stamford: Cengage Learning. 10th edition. Chapters 13 (pp. 342–378), 15 (pp. 405–441).



Outline

- **Introduction**
 - Bivariate regression
 - Multivariate regression
 - Standardized coefficients (b^*)
 - Statistical significance (t -test)
 - Multiple correlation (R^2)
 - Assumptions: Gauss-Markov theorem
- **Meaning of linear regression**
 - Example: Income = F(age, education)
- **Determining normality**
 - Example: $\ln(\text{income}) = F(\text{age}, \text{education})$
- **Predicted values**
- **Residual analysis with graphs**
 - Example: OLS with age and age squared
- **Dummy variables**
 - Example: Full OLS model



Introduction

- Ordinary least squares (OLS) regression (linear regression)
 - Important technique to estimate associations of several independent variables (x_1, x_2, \dots, x_k) with a dependent variable (y) at the interval-ratio level of measurement
 - Variables are at the interval-ratio level, but we can include ordinal and nominal variables as dummy variables
 - Each independent variable has a linear relationship with the dependent variable
 - Independent variables are uncorrelated with each other
 - When these and other requirements are violated (as they often are), this technique will produce biased and/or inefficient estimates



Correlation vs. causation

- Correlation and causation are different
 - Strong associations (correlation) may be used as evidence of causal relationships (causation)
 - Associations do not prove variables are causally related
- We might have problems of reverse causality
 - e.g., immigration increases competition in the labor market and affects earnings
 - Availability of jobs and income levels influence migration

Migration  **Earnings**



Bivariate and multivariate models

- Bivariate (simple) regression equation


$$y = a + bx = \beta_0 + \beta_1x$$

- $a = \beta_0 = y$ intercept (constant)
- $b = \beta_1 =$ slope

- Multivariate (multiple) regression equation

$$y = a + b_1x_1 + b_2x_2 = \beta_0 + \beta_1x_1 + \beta_2x_2$$

- $b_1 = \beta_1 =$ partial slope of the linear relationship between the first independent variable (x_1) and y
- $b_2 = \beta_2 =$ partial slope of the linear relationship between the second independent variable (x_2) and y



TEXAS A&M
UNIVERSITY.


Bivariate regression

$$y = a + bx = \beta_0 + \beta_1x$$

- $a = \beta_0 = y$ intercept (constant)
- $b = \beta_1 =$ slope
- In a scatterplot
 - The independent variable (x) is displayed along the horizontal axis
 - The dependent variable (y) is displayed along the vertical axis
 - Each dot on a scatterplot is a case
 - The dot is placed at the intersection of the case's scores on x and y

Example of a scatterplot

- Number of children (x) and hours per week husband spends on housework (y) at dual-career households




Regression line

- A regression line is added to the graph
- It summarizes the linear correlation between x and y
 - This straight line connects all of the dots
 - Or this line comes as close as possible to connecting all of the dots

Scatterplot with regression line

Husband's Housework by Number of Children




Prediction

- Scatterplots can be used to predict values of Y (Y' or \hat{Y}) based on values of X
- Locate a particular X value on the horizontal axis
- Draw a vertical line up to the regression line
- Then draw a horizontal line over to the vertical axis

Example of prediction

Predicting Husband's Housework



Estimating the regression line


- The regression line touches each conditional mean of Y
 - Or the line comes as close as possible to all scores
- The dots above each value of X can be thought of as conditional distributions of Y
 - In previous chapters, column percentages were the conditional distributions of Y for each value of X



Conditional means of Y


- Conditional means of Y are found by summing all Y values for each value of X and dividing by the number of cases

Original data



Conditional means of Y

Conditional means of Y



Number of Children (X)	Husband's Housework (Y)	Conditional Mean of Y
1	1, 2, 3, 5	2.75
2	3, 1	2.00
3	5, 0	2.50
4	6, 3	4.50
5	7, 4	5.50



Estimating coefficients

- Ordinary least squares (OLS) simple regression
 - OLS: linear regression
 - Simple: only one independent variable

$$Y = a + bX = \beta_0 + \beta_1 X$$

- Where
 - Y = score on the dependent variable
 - X = score on the independent variable
 - $a = \beta_0$ = the Y intercept or the point where the regression line crosses the Y axis
 - $b = \beta_1$ = slope of the regression line or the amount of change produced in Y by a unit change in X



Computing the slope (b)

- Before using the formula for the regression line, we need to estimate a and b
- First, estimate b

$$b = \frac{\sum(X - \bar{X})(Y - \bar{Y})}{\sum(X - \bar{X})^2}$$

- The numerator of the formula is the “covariation” of X and Y
 - How much X and Y vary together
 - Its value reflects the direction and strength of the association between X and Y



Computing the Y intercept (a)

- The intercept (a) is the point where the regression line crosses the Y axis
- Estimate a using the mean for X, the mean for Y, and b

$$a = \bar{Y} - b\bar{X}$$

Example

- Number of children (X) and hours per week husband spends on housework (Y) at dual-career households

Number of Children and Husband's Contribution to Housework
(fictitious data)

Family	Number of Children	Hours per Week Husband Spends on Housework
A	1	1
B	1	2
C	1	3
D	1	5
E	2	3
F	2	1
G	3	5
H	3	0
I	4	6
J	4	3
K	5	7
L	5	4



Example: calculation table

- Calculation of b is simplified if you set up a computation table

Computation of the Slope (b)

1	2	3	4	5	6
X	$X - \bar{X}$	Y	$Y - \bar{Y}$	$(X - \bar{X})(Y - \bar{Y})$	$(X - \bar{X})^2$
1	-1.67	1	-2.33	3.89	2.79
1	-1.67	2	-1.33	2.22	2.79
1	-1.67	3	-0.33	0.55	2.79
1	-1.67	5	1.67	-2.79	2.79
2	-0.67	3	-0.33	0.22	0.45
2	-0.67	1	-2.33	1.56	0.45
3	0.33	5	1.67	0.55	0.11
3	0.33	0	-3.33	-1.10	0.11
4	1.33	6	2.67	3.55	1.77
4	1.33	3	-0.33	-0.44	1.77
5	2.33	7	3.67	8.55	5.43
5	2.33	4	0.67	1.56	5.43
<u>32</u>	<u>-0.04</u>	<u>40</u>	<u>0.04</u>	<u>18.32</u>	<u>26.68</u>

$$\bar{X} = \frac{32}{12} = 2.67$$

$$\bar{Y} = \frac{40}{12} = 3.33$$



Example: slope and intercept

- Based on previous table, estimate the slope (b)

$$b = \frac{\sum(X - \bar{X})(Y - \bar{Y})}{\sum(X - \bar{X})^2} = \frac{18.32}{26.68} = 0.69$$

- Estimate the intercept (a)

$$a = \bar{Y} - b\bar{X} = 3.33 - (0.69)(2.67) = 1.49$$

Example: interpretations

- Regression equation with $a=1.49$ and $b=0.69$

$$Y' = 1.49 + (0.69)X$$

– $b = 0.69$

- For every additional child in the dual-career household, husbands perform on average an additional 0.69 hours (around 36 minutes) of housework per week


– $a = 1.49$

- The regression line crosses the Y axis at 1.49
- When there are zero children in a dual-career household, husbands perform on average 1.49 hours of housework per week



Example: coefficients

Husband's Housework by Number of Children



Example: predictions

- What is the predicted value of Y (Y') when X equals 6?
$$Y' = 1.49 + (0.69)X = 1.49 + (0.69)(6) = 5.63$$
 - In dual-career families with 6 children, the husband is predicted to perform on average 5.63 hours of housework a week
- What about when X equals 7?
$$Y' = 1.49 + (0.69)X = 1.49 + (0.69)(7) = 6.32$$
 - In dual-career families with 7 children, the husband is predicted to perform on average 6.32 hours of housework a week
 - Notice how the difference in these two predicted values equals b ($6.32 - 5.63 = 0.69$)



GSS: Income = F(Education)

```

***Dependent variable: Respondent's income (conrinc)
***Independent variable: Years of schooling (educ)

***Scatterplot with regression line
twoway scatter conrinc educ || lfit conrinc educ, ytitle(Respondent's income) xtitle(Years of schooling)

***Regression coefficients
***Least-squares regression model
***They can be reported in the footnote of the scatterplot
svy: reg conrinc educ

```

```

. svy: reg conrinc educ
(running regress on estimation sample)

```

Survey: Linear regression


Number of strata	=	65	Number of obs	=	1,631
Number of PSUs	=	130	Population size	=	1,694.7478
			Design df	=	65
			F(1, 65)	=	88.15
			Prob > F	=	0.0000
			R-squared	=	0.1147

conrinc	Linearized				
	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]
educ	4326.103	460.7631	9.39	0.000	3405.896 5246.311
_cons	-26219.18	5819.513	-4.51	0.000	-37841.55 -14596.81



Income by education

Figure 1. Respondent's income by years of schooling, U.S. adult population, 2016



$$\text{Income} = -26,219.18 + 4,326.10(\text{Years of schooling})$$

Note: The scatterplot was generated without the complex survey design of the General Social Survey. The regression was generated taking into account the complex survey design of the General Social Survey.

Source: 2016 General Social Survey.

ACS: Income = F(Age)

***Dependent variable: Wage and salary income (income)

***Independent variable: Age (age)

***Scatterplot with regression line

twoway (scatter income age) (lfit income age) if income!=0, ytitle(Wage and salary income) xtitle(Age)

```
. svy, subpop(if income!=. & income!=0): reg income age
(running regress on estimation sample)
```

Survey: Linear regression

Number of strata = 2,351
Number of PSUs = 1,410,976


Number of obs = 3,214,539
Population size = 327,167,439
Subpop. no. obs = 1,574,313
Subpop. size = 163,349,075
Design df = 1,408,625
F(1,1408625) = 57648.04
Prob > F = 0.0000
R-squared = 0.0449

income	Linearized					
	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
age	888.2282	3.699409	240.10	0.000	880.9775	895.479
_cons	13447.38	138.3572	97.19	0.000	13176.21	13718.56



Income by age


Figure 1. Wage and salary income by age, U.S. 2018



$$\text{Income} = 13,447.38 + 888.23(\text{Age})$$

Note: The scatterplot was generated without the ACS complex survey design. The regression was generated taking into account the ACS complex survey design. Only people with some wage and salary income are included.

Source: 2018 American Community Survey (ACS).



TEXAS A&M
UNIVERSITY.

Multivariate regression

$$y = a + b_1x_1 + b_2x_2 = \beta_0 + \beta_1x_1 + \beta_2x_2$$

- $a = \beta_0$ = the y intercept (constant), where the regression line crosses the y axis
- $b_1 = \beta_1$ = partial slope for x_1 on y
 - β_1 indicates the change in y for one unit change in x_1 , controlling for x_2
- $b_2 = \beta_2$ = partial slope for x_2 on y
 - β_2 indicates the change in y for one unit change in x_2 , controlling for x_1


Partial slopes (β)

- The partial slopes (β) indicate the effect of each independent variable on y
- While controlling for the effect of the other independent variables
- This control is called *ceteris paribus*
 - Other things equal
 - Other things held constant
 - All other things being equal




Ceteris paribus

$$\text{Income} = \beta_0 + \beta_1 \text{education} + \beta_2 \text{experience} + e$$



Ceteris paribus

$$\text{Income} = \beta_0 + \beta_1 \text{education} + \beta_2 \text{experience} + e$$




Ceteris paribus

$$\text{Income} = \beta_0 + \beta_1 \text{education} + \beta_2 \text{experience} + e$$

Low
education

High
education




These two effects
will result on

β_2



Ceteris paribus

$$\text{Income} = \beta_0 + \beta_1 \text{education} + \beta_2 \text{experience} + e$$



Interpretation of partial slopes

- The partial slopes show the effects of the independent variables (x_1, x_2) in their original units
- These values can be used to predict scores on the dependent variable (y)
- Partial slopes must be computed before computing the y intercept (β_0)



Formulas of partial slopes

$$b_1 = \beta_1 = \left(\frac{s_y}{s_1} \right) \left(\frac{r_{y1} - r_{y2}r_{12}}{1 - r_{12}^2} \right)$$

$$b_2 = \beta_2 = \left(\frac{s_y}{s_2} \right) \left(\frac{r_{y2} - r_{y1}r_{12}}{1 - r_{12}^2} \right)$$

$b_1 = \beta_1$ = partial slope of x_1 on y

$b_2 = \beta_2$ = partial slope of x_2 on y

s_y = standard deviation of y

s_1 = standard deviation of the first independent variable (x_1)

s_2 = standard deviation of the second independent variable (x_2)

r_{y1} = bivariate correlation between y and x_1

r_{y2} = bivariate correlation between y and x_2

r_{12} = bivariate correlation between x_1 and x_2



Formula of constant

- Once b_1 (β_1) and b_2 (β_2) have been calculated, use those values to calculate the y intercept (β_0)

$$a = \bar{y} - b_1 \bar{x}_1 - b_2 \bar{x}_2$$

$$\beta_0 = \bar{y} - \beta_1 \bar{x}_1 - \beta_2 \bar{x}_2$$

Income = F(age, education)

```
. ***No weights
. reg income age educgr
```

Source	SS	df	MS	Number of obs	=	127,785
Model	8.2170e+13	2	4.1085e+13	F(2, 127782)	=	11557.33
Residual	4.5425e+14	127,782	3.5549e+09	Prob > F	=	0.0000
Total	5.3642e+14	127,784	4.1979e+09	R-squared	=	0.1532
				Adj R-squared	=	0.1532
				Root MSE	=	59623

income	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
age	724.3054	11.11857	65.14	0.000	702.5132	746.0976
educgr	18177.19	140.4437	129.43	0.000	17901.92	18452.45
_cons	-32363.61	614.972	-52.63	0.000	-33568.95	-31158.28



Summary of Stata weights

WEIGHTS IN FREQUENCY DISTRIBUTIONS

Weight unit of measurement	Expand to population size	Maintain sample size
Discrete	fweight	aweight
Continuous	iweight	

WEIGHTS IN STATISTICAL REGRESSIONS should maintain sample size

Robust standard error	Adjusted R ² , TSS, ESS, RSS
pweight	aweight
reg y x, vce(robust) reg y x, vce(cluster area)	outreg2



Example: Coefficients (β)


```
. ***Complex survey design
. svyset cluster [pweight=perwt], strata(strata)

. ***Use complex survey design
. svy: reg income age educgr
(running regress on estimation sample)
```

Survey: Linear regression

```
Number of strata   =          212
Number of PSUs    =       79,499
Number of obs     =      127,785
Population size   =  13,849,398
Design df        =       79,287
F( 2, 79286)     =      5751.26
Prob > F         =          0.0000
R-squared        =          0.1652
```

	Coef.	Linearized Std. Err.	t	P> t	[95% Conf. Interval]	
income						
age	796.3443	11.73077	67.89	0.000	773.3521	819.3366
educgr	16863.33	179.705	93.84	0.000	16511.11	17215.55
_cons	-31880.99	661.937	-48.16	0.000	-33178.38	-30583.59



TEXAS A&M
UNIVERSITY.

Standardized coefficients (b^*)

- Partial slopes ($b_1=\beta_1$; $b_2=\beta_2$) are in the original units of the independent variables
 - This makes assessing relative effects of independent variables difficult when they have different units
 - It is easier to compare if we standardize to a common unit by converting to Z scores
- Compute beta-weights (b^*) to compare relative effects of the independent variables
 - Amount of change in the standardized scores of y for a one-unit change in the standardized scores of each independent variable
 - While controlling for the effects of all other independent variables
 - They show the amount of change in standard deviations in y for a change of one standard deviation in each x



Formulas

- Formulas for standardized coefficients

$$b_1^* = b_1 \left(\frac{s_1}{s_y} \right) = \beta_1^* = \beta_1 \left(\frac{s_1}{s_y} \right)$$

$$b_2^* = b_2 \left(\frac{s_2}{s_y} \right) = \beta_2^* = \beta_2 \left(\frac{s_2}{s_y} \right)$$

Standardized coefficients

- Standardized regression equation

$$Z_y = a_z + b_1^*Z_1 + b_2^*Z_2$$

- Z indicates that all scores have been standardized to the normal curve

$$Z_i = \frac{x_i - \bar{x}}{s}$$

- The y intercept will always equal zero once the equation is standardized

$$Z_y = b_1^*Z_1 + b_2^*Z_2$$




Example: Standardized beta (b^*)

```
. ***Standardized regression coefficients
. ***(i.e., standardized partial slopes, beta-weights)
. ***It does not allow the use of complex survey design
. ***Use pweight to maintain sample size and estimate robust standard errors
. reg income age educgr [pweight=perwt], beta
(sum of wgt is 13,849,398)
```

```
Linear regression                               Number of obs       =      127,785
                                                F(2, 127782)       =      5873.56
                                                Prob > F            =      0.0000
                                                R-squared           =      0.1652
                                                Root MSE           =      54147
```

income	Coef.	Robust Std. Err.	t	P> t	Beta
age	796.3443	11.46129	69.48	0.000	.1943233
educgr	16863.33	177.6256	94.94	0.000	.3368842
_cons	-31880.99	649.8899	-49.06	0.000	.





TEXAS A&M
UNIVERSITY.

Statistical significance (*t*-test)

- In a simple linear regression, the test of statistical significance for a β coefficient (*t*-test) is estimated as

$$t = \frac{\hat{\beta}}{SE_{\hat{\beta}}} = \frac{\hat{\beta}}{\sqrt{\frac{MSE}{S_{xx}}}} = \frac{\hat{\beta}}{\sqrt{\frac{RSS}{df * S_{xx}}}} = \frac{\hat{\beta}}{\sqrt{\frac{\sum_i (y_i - \hat{y}_i)^2}{(n - 2) \sum_i (x_i - \bar{x})^2}}}$$

- $SE_{\hat{\beta}}$: standard error of β
- MSE : mean squared error = RSS / df
- RSS : residual sum of squares = $\sum_i (y_i - \hat{y}_i)^2 = \sum_i \hat{e}_i^2$
- df : degrees of freedom = $n-2$ for simple linear regression
 - 2 statistics (slope and intercept) are estimated to calculate sum of squares
- S_{xx} : corrected sum of squares for x (total sum of squares)



Statistical power


- Statistical power for regression analysis is the probability of finding a significant coefficient ($\hat{\beta} \neq 0$), when there is a significant relationship in the population ($\beta \neq 0$)
 - Power is dependent on the confidence level, size of coefficient (magnitude), and sample size
 - Small samples might not capture enough variation among observations
 - If we have large samples, we tend to have statistical significance (as measured by t -test), even for coefficients ($\hat{\beta}$) with small magnitude

$$\uparrow t = \frac{\hat{\beta}}{SE_{\hat{\beta}}} = \frac{\hat{\beta}}{\sqrt{\frac{MSE}{S_{xx}}}} = \frac{\hat{\beta}}{\sqrt{\frac{RSS}{df * S_{xx}}}} = \frac{\hat{\beta}}{\sqrt{\frac{\sum_i (y_i - \hat{y}_i)^2}{(n - 2) \sum_i (x_i - \bar{x})^2}}}$$

↑ ↓

t distribution ($df = 2$)

- **Bigger the t -test**
 - Stronger the statistical significance
- **Smaller the p -value**
 - Smaller the probability of not rejecting the null hypothesis
 - Tend to accept alternative (research) hypothesis



Decisions about hypotheses

Hypotheses	$p < \alpha$	$p > \alpha$
Null hypothesis (H_0)	Reject	Do not reject
Alternative hypothesis (H_1)	Accept	Do not accept

– ***p*-value** is the probability of not rejecting the null hypothesis

– If a statistical software gives only the two-tailed *p*-value, divide it by 2 to obtain the one-tailed *p*-value

Significance level (α)	Confidence level (success rate)
0.10 (10%)	90%
0.05 (5%)	95%
0.01 (1%)	99%
0.001 (0.1%)	99.9%




Example: Statistical significance

```
. ***Use complex survey design  
. svy: reg income age educgr  
(running regress on estimation sample)
```

Survey: Linear regression

```
Number of strata   =      212           Number of obs     =    127,785  
Number of PSUs    =    79,499         Population size   =  13,849,398  
                                                           Design df        =    79,287  
                                                           F( 2, 79286)    =    5751.26  
                                                           Prob > F        =    0.0000  
                                                           R-squared       =    0.1652
```

	Coef.	Linearized Std. Err.	t	P> t	[95% Conf. Interval]	
income						
age	796.3443	11.73077	67.89	0.000	773.3521	819.3366
educgr	16863.33	179.705	93.84	0.000	16511.11	17215.55
_cons	-31880.99	661.937	-48.16	0.000	-33178.38	-30583.59



TEXAS A&M
UNIVERSITY.

Multiple correlation (R^2)

- The coefficient of multiple determination (R^2) measures how much of the dependent variable (y) is explained by all independent variables ($x_1, x_2, x_3, \dots, x_k$) combined
- R^2 is an estimation of the percentage of the variation in y that is explained by variations in all independent variables in the population
- The coefficient of multiple determination is an indicator of the strength of the entire regression equation



R^2 estimation

- For a regression with two independent variables, this is the equation to estimate R^2

$$R^2 = r_{y1}^2 + r_{y2.1}^2(1 - r_{y1}^2)$$

- R^2 = coefficient of multiple determination
- r_{y1}^2 = coefficient of determination for y and x_1 (or amount of variation in y explained by x_1)
- $r_{y2.1}^2$ = partial correlation of y and x_2 , while controlling for x_1 (or amount of variation in y explained by x_2 , after x_1 is controlled)
- $(1 - r_{y1}^2)$ = amount of variation remaining in y , after controlling for x_1

Partial correlation of y and x_2

- Before estimating R^2 , we need to estimate the partial correlation of y and x_2 , while controlling for x_1 ($r_{y2.1}$)

$$r_{y2.1} = \frac{r_{y2} - (r_{y1})(r_{12})}{\sqrt{1 - r_{y1}^2} \sqrt{1 - r_{12}^2}}$$

- We need three correlations
 - Bivariate correlation between y and x_1 (r_{y1})
 - Bivariate correlation between y and x_2 (r_{y2})
 - Bivariate correlation between x_1 and x_2 (r_{12})

Explaining R^2 estimation

$$R^2 = r_{y1}^2 + r_{y2.1}^2(1 - r_{y1}^2)$$

- If the partial correlation of y and x_2 , while controlling for x_1 ($r_{y2.1}$), is not equal to zero
 - R^2 will necessarily increase by adding x_2
 - Any variable x will have a non-zero correlation with y
 - In real databases, y and any x don't have correlation exactly equal to zero
- Thus, more independent variables (even if not related to theory) will generate higher R^2

R^2 and independent variables

- Selection of independent variables based on R^2 size might generate unreasonable models
- There is nothing in the hypotheses of linear models that require a minimum value for R^2
- Models with small R^2 might mean that we didn't include important independent variables
 - It doesn't mean necessarily that non-observed factors (residuals) are correlated with independent variables
- R^2 size doesn't have influence on the mean of residuals being equal to zero

R^2 in terms of variance

- R^2 can also be written in terms of variance of y in the population (σ_y^2) and variance of error term (residual u) in the population (σ_u^2)

$$R^2 = 1 - \sigma_u^2 / \sigma_y^2$$

- R^2 is the proportion of variation in y explained by all independent variables...

$TSS = ESS + RSS$

$$\sum_{i=1}^n (y_i - \bar{y})^2 = \sum_{i=1}^n (\hat{y}_i - \bar{y})^2 + \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

– Total sum of squares (TSS)

- Sum of squares total (SST)
- $TSS = \sum_{i=1}^n (y_i - \bar{y})^2$
- df (degrees of freedom) = $n-1$, where n is the sample size
- Average total sum of squares = $TSS / df = TSS / (n-1)$

– Explained sum of squares (ESS)

- Sum of squares due to regression (SSR), model sum of squares (MSS)
- $ESS = \sum_{i=1}^n (\hat{y}_i - \bar{y})^2$
- df = k , where k is the number of independent variables
- Average explained sum of squares = $ESS / df = ESS / k$

– Residual sum of squares (RSS)

- Sum of squared errors of prediction (SSE)
- $RSS = \sum_{i=1}^n (y_i - \hat{y}_i)^2 = \sum_{i=1}^n \hat{e}_i^2$
- df = $n-k-1$
- Average residual sum of squares = $RSS / df = RSS / (n-k-1)$



R^2 in terms of variance

- Total sum of squares equal explained sum of squares plus residual sum of squares

$$TSS = ESS + RSS$$

$$TSS/TSS = (ESS + RSS)/TSS$$

$$1 = ESS/TSS + RSS/TSS$$

$$ESS/TSS = 1 - RSS/TSS$$

- R^2 is the proportion of variation in y explained by all independent variables

$$R^2 = ESS / TSS$$

$$R^2 = 1 - RSS / TSS$$

$$R^2 = 1 - (RSS/n) / (TSS/n)$$

$$R^2 = 1 - \sigma_u^2 / \sigma_y^2$$



Adjusted R^2

- We can replace RSS/n and TSS/n by non-biased terms for σ_u^2 and σ_y^2

$$\text{Adjusted } R^2 = 1 - [RSS/(n-k-1)] / [TSS/(n-1)]$$

- Adjusted R^2 doesn't correct for possible bias of R^2 estimating the true population R^2
- But it penalizes for the inclusion of redundant independent variables
- k is the number of independent variables
- Negative adjusted R^2 indicates a poor overall fit

$$\downarrow \text{Adjusted } R^2 = 1 - \frac{1 - R^2}{n - \uparrow k - 1} \uparrow$$



Comparing models

- We can compare adjusted R^2 of models with different forms of independent variables

$$y = \beta_0 + \beta_1 \log(x) + u$$

$$y = \beta_0 + \beta_1 x + \beta_2 x^2 + u$$

- We cannot use R^2 or adjusted R^2 to choose between different forms of dependent variable
- Different forms of y have different amounts of variation to be explained



Example: R^2 , Adjusted R^2


```
. ***Use aweight to estimate adjusted R-squared
. ***pweight and complex survey design omit sum of squares and adjusted R-squared
. reg income age educgr [aweight=perwt]
(sum of wgt is 13,849,398)
```

Source	SS	df	MS
Model	7.4126e+13	2	3.7063e+13
Residual	3.7465e+14	127,782	2.9319e+09
Total	4.4877e+14	127,784	3.5120e+09

```
Number of obs   = 127,785
F(2, 127782)    = 12641.17
Prob > F        = 0.0000
R-squared       = 0.1652
Adj R-squared   = 0.1652
Root MSE       = 54147
```

income	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
age	796.3443	10.53436	75.59	0.000	775.6972	816.9915
educgr	16863.33	128.6752	131.05	0.000	16611.13	17115.53
_cons	-31880.99	554.2213	-57.52	0.000	-32967.25	-30794.72





TEXAS A&M
UNIVERSITY.

Gauss-Markov theorem

- The Gauss-Markov theorem states that if the linear regression model satisfies classical assumptions
 - Then ordinary least squares (OLS) regression produces unbiased estimates that have the smallest variance of all possible linear estimators
 - We should have a random sample of n observations for the population model
 - *Best Linear Unbiased Estimators (BLUEs)*



1. Linear in parameters


- The regression model is linear in the coefficients and the error term
 - An increase of one unit in an independent variable makes the expected value of y to vary by the magnitude of the correspondent β
 - All terms in the model are either the constant or a parameter multiplied by an independent variable
 - The population model can be written as
$$y = \beta_0 + \beta_1x_1 + \beta_2x_2 + \beta_3x_3 + \dots + \beta_kx_k + e$$
 - $\beta_0, \beta_1, \dots, \beta_k$ represent unknown parameters
 - Error term is known as the residual ($e, \epsilon, \text{ or } u$)
 - It is an unobserved random error
 - It is the variation in y that the model doesn't explain



Conditional means of y

- For any value of x , the distribution of y is centered around the expected value of y given x

$\hat{y} = E(y|x)$ as a linear function of x



2. No perfect collinearity

- No independent variable is a perfect linear function of other independent variables
 - No independent variable is constant and there are no exact linear relations among independent variables
- Independent variables should be associated among themselves, but there should be **no perfect collinearity**
 - e.g., one variable should not be the multiple of another one
- High levels of correlation among independent variables and small sample size increase standard errors of β
 - This decreases statistical significance: $t = \beta / SE_{\beta}$
- High correlation (but not perfect) among independent variables is not desirable (**multicollinearity**)



3. All x are uncorrelated with e

- All independent variables (x) are uncorrelated with the error term (e)
 - If an independent variable is correlated with the error term, the independent variable can be used to predict the error term
 - This violates the notion that the error term represents unpredictable random error
- This assumption is referred to as exogeneity
 - When this type of correlation exists, there is endogeneity
 - There is reverse causality between independent and dependent variables, omitted variable bias, or measurement error




4. Uncorrelated observations of e

- Observations of the error term (e) are uncorrelated with each other
 - One observation of the error term should not predict the next observation


- Verify by graphing the residuals in the order that the data was collected
 - We want to see randomness in the plot

E.g. observations of e are correlated



5. Error term has mean of zero


- The error term has as population mean of zero
 - The expected value (mean) of the unobserved random error (e) is zero, given any values of the independent variables
 - $E(e|x_1, x_2, \dots, x_k) = 0$
- Residuals = $e = y_i - \hat{y}_i$
 - Observed minus fitted
 - Observed minus predicted
 - Sum of residuals (population mean) should be zero




6. Homoscedasticity

- The error term has a constant variance (no heteroscedasticity)
 - Variance of errors (e) should be consistent for all observations
 - Variance does not change for each observation or range of observations
 - If this assumption is violated, the model has heteroscedasticity

Homoscedasticity




Heteroscedasticity




7. Optional: e is normally distributed

- The error term (e) should be normally distributed
 - OLS does not require that the error term follows a normal distribution to produce unbiased estimates with minimum variance
 - But satisfying this assumption allows us to perform statistical hypothesis testing and generate reliable confidence and prediction intervals

E.g. residuals are normally distributed





TEXAS A&M
UNIVERSITY.

Meaning of linear regression

- Ordinary least squares regression is commonly named linear regression
- The model is linear in the parameters: β_0, β_1, \dots
- An increase of one unit in an independent variable makes the expected value of y to vary by the magnitude of the correspondent β
- However, it allows us to include non-linear associations

No restrictions

- There are no restrictions of how y and x are associated with the original dependent and independent variables
- We can use natural logarithm, squared values, squared root, dummy independent variables...
- The **interpretation** of coefficients depends of how y and x are estimated and included in the regression



Interpretation of coefficients

- An increase of one unit in x increases y by β_1 units

$$y = \beta_0 + \beta_1 x + e$$

- An increase of 1% in x increases y by $(\beta_1/100)$ units

$$y = \beta_0 + \beta_1 \log(x) + e$$

- An increase of one unit in x increases y by $(100*\beta_1)\%$
 - Exact percentual change with semi-elasticity $\{[\exp(\beta_1) - 1]*100\}$

$$\log(y) = \beta_0 + \beta_1 x + e$$

- An increase of 1% in x increases y by $\beta_1\%$
 - Constant elasticity model
 - Elasticity is the ratio of the percentage change in y to the percentage change in x

$$\log(y) = \beta_0 + \beta_1 \log(x) + e$$




Logarithm functional forms

Model	Dependent variable	Independent variable	Interpretation of β_1
linear	y	x	$\Delta y = \beta_1 \Delta x$
linear-log	y	log(x)	$\Delta y = (\beta_1 / 100) \% \Delta x$
log-linear (semi-log)	log(y)	x	$\% \Delta y = (100 \beta_1) \Delta x$
log-log	log(y)	log(x)	$\% \Delta y = \beta_1 \% \Delta x$




Linear

x	y
1	1
2	2
3	3
4	4
5	5
6	6
7	7
8	8
9	9
10	10




x	y
1	10
2	9
3	8
4	7
5	6
6	5
7	4
8	3
9	2
10	1




Linear-Log

log(x)	y
0.0	1
0.7	2
1.1	3
1.4	4
1.6	5
1.8	6
1.9	7
2.1	8
2.2	9
2.3	10




log(x)	y
0.0	10
0.7	9
1.1	8
1.4	7
1.6	6
1.8	5
1.9	4
2.1	3
2.2	2
2.3	1




Log-Linear

x	log(y)
1	0.0
2	0.7
3	1.1
4	1.4
5	1.6
6	1.8
7	1.9
8	2.1
9	2.2
10	2.3




x	log(y)
1	2.3
2	2.2
3	2.1
4	1.9
5	1.8
6	1.6
7	1.4
8	1.1
9	0.7
10	0.0





Log-Log

log(x)	log(y)
0.0	0.0
0.7	0.7
1.1	1.1
1.4	1.4
1.6	1.6
1.8	1.8
1.9	1.9
2.1	2.1
2.2	2.2
2.3	2.3



log(x)	log(y)
0.0	2.3
0.7	2.2
1.1	2.1
1.4	1.9
1.6	1.8
1.8	1.6
1.9	1.4
2.1	1.1
2.2	0.7
2.3	0.0





TEXAS A&M
UNIVERSITY.

Income = F(age, education)

```
. ***Use complex survey design
. svy: reg income age educgr
(running regress on estimation sample)
```

Survey: Linear regression

```
Number of strata   =      212
Number of PSUs    =    79,499
Number of obs     =    127,785
Population size   =  13,849,398
Design df        =      79,287
F( 2, 79286)     =    5751.26
Prob > F         =      0.0000
R-squared        =      0.1652
```

income	Coef.	Linearized Std. Err.	t	P> t	[95% Conf. Interval]	
age	796.3443	11.73077	67.89	0.000	773.3521	819.3366
educgr	16863.33	179.705	93.84	0.000	16511.11	17215.55
_cons	-31880.99	661.937	-48.16	0.000	-33178.38	-30583.59

Interpretation of coefficients

(income with continuous independent variables)

- Coefficient for **age** equals 796.34
 - When age increases by one unit, income increases on average by **796.34 dollars**, controlling for education
- Coefficient for **education** equals 16,863.33
 - When education increases by one unit, income increases on average by **16,863.33 dollars**, controlling for age



Standardized coefficients

```
. ***Standardized regression coefficients
. ***(i.e., standardized partial slopes, beta-weights)
. ***It does not allow the use of complex survey design
. ***Use pweight to maintain sample size and estimate robust standard errors
. reg income age educgr [pweight=perwt], beta
(sum of wgt is 13,849,398)
```

```
Linear regression                Number of obs    =    127,785
                                F(2, 127782)    =    5873.56
                                Prob > F             =    0.0000
                                R-squared            =    0.1652
                                Root MSE         =    54147
```

income	Coef.	Robust Std. Err.	t	P> t	Beta
age	796.3443	11.46129	69.48	0.000	.1943233
educgr	16863.33	177.6256	94.94	0.000	.3368842
_cons	-31880.99	649.8899	-49.06	0.000	.

Interpretation of standardized

(income with continuous independent variables)

- Coefficient for **age** equals 0.1943
 - When age increases by one standard deviation, income increases on average by **0.1943 standard deviations**, controlling for education
- Coefficient for **education** equals 0.3369
 - When education increases by one standard deviation, income increases on average by **0.3369 standard deviations**, controlling for age

Adjusted R^2

```
. ***Use aweight to estimate adjusted R-squared
. ***pweight and complex survey design omit sum of squares and adjusted R-squared
. reg income age educgr [aweight=perwt]
(sum of wgt is 13,849,398)
```

Source	SS	df	MS	Number of obs	=	127,785
Model	7.4126e+13	2	3.7063e+13	F(2, 127782)	=	12641.17
Residual	3.7465e+14	127,782	2.9319e+09	Prob > F	=	0.0000
				R-squared	=	0.1652
				Adj R-squared	=	0.1652
Total	4.4877e+14	127,784	3.5120e+09	Root MSE	=	54147

income	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
age	796.3443	10.53436	75.59	0.000	775.6972	816.9915
educgr	16863.33	128.6752	131.05	0.000	16611.13	17115.53
_cons	-31880.99	554.2213	-57.52	0.000	-32967.25	-30794.72




TEXAS A&M
UNIVERSITY.

Determining normality


- Some statistical methods require random selection of respondents from a population with normal distribution for its variables
 - OLS regressions require normal distribution for its interval-ratio-level variables
 - We can analyze histograms, boxplots, outliers, quantile-normal plots, and measures of skewness and kurtosis to determine if variables have a normal distribution



Histogram of income



Boxplot of income



Source: 2018 American Community Survey.




Quantile-normal plots


- A quantile-normal plot is a scatter plot
 - One axis has quantiles of the original data
 - The other axis has quantiles of the normal distribution
- If the points do not form a straight line or if the points have a non-linear symmetric pattern
 - The variable does not have a normal distribution
- If the pattern of points is roughly straight
 - The variable has a distribution close to normal
- If the variable has a normal distribution
 - The points would exactly overlap the diagonal line




Quantile-normal plots reflect distribution shapes




Heavy Tails, High and Low Outliers




Light Tails, No Outliers




Positive Skew, High Outliers



Negative Skew, Low Outliers




Granularity
(discrete values)



Two Peaks, Central Gap
(bimodal)

Quantile-normal plot of income



Source: 2018 American Community Survey.

Skewness

- Skewness is a measure of symmetry
 - A distribution is symmetric if it looks the same to the left and right of the center point
 - Skewness for a normal distribution is zero
 - Negative values for the skewness indicate variable is skewed to the left (left tail is long relative to the right tail)
 - Positive values for the skewness indicate variable is skewed to the right (right tail is long relative to the left tail)
- Rule of thumb
 - Skewness between -0.5 and 0.5 : variable is fairly symmetrical
 - Skewness between -1 and -0.5 or between 0.5 and 1 : variable moderately skewed
 - Skewness less than -1 or greater than 1 : variable is highly skewed

Kurtosis

- Kurtosis is a measure of whether the data are heavy-tailed or light-tailed relative to a normal distribution
 - Variables with high kurtosis tend to have heavy tails or outliers
 - Variables with low kurtosis tend to have light tails or lack of outliers
 - A uniform distribution would be the extreme case
 - **The kurtosis for a standard normal distribution is three**
- Excess kurtosis
 - Some sources subtract 3 from the kurtosis
 - The standard normal distribution has an excess kurtosis of zero
 - Positive excess kurtosis indicates a “heavy-tailed” distribution
 - Negative excess kurtosis indicates a “light tailed” distribution

Source: <https://www.itl.nist.gov/div898/handbook/eda/section3/eda35b.htm>
<https://www.spcforexcel.com/knowledge/basic-statistics/are-skewness-and-kurtosis-useful-statistics>
<https://www.stata-journal.com/sjpdf.html?articlenum=st0204>



Skewness and Kurtosis

```
. sum income if income!=0 [fweight=perwt], d
```

income

Percentiles		Smallest		
1%	500	4		
5%	2400	4		
10%	5600	4	Obs	13,849,398
25%	16000	4	Sum of Wgt.	13,849,398
50%	34000		Mean	48713.66
		Largest	Std. Dev.	59261.63
75%	60000	468000		
90%	100000	468000	Variance	3.51e+09
95%	136000	468000	Skewness	4.20286
99%	468000	468000	Kurtosis	27.61478

Power transformation

- Lawrence Hamilton (“Regression with Graphics”, 1992, p.18–19)

$$y^3 \rightarrow q = 3$$

$$y^2 \rightarrow q = 2$$

$$y^1 \rightarrow q = 1$$

$$y^{0.5} \rightarrow q = 0.5$$

$$\log(y) \rightarrow q = 0$$


$$-(y^{-0.5}) \rightarrow q = -0.5$$

$$-(y^{-1}) \rightarrow q = -1$$


- $q > 1$: reduce concentration on the right (reduce negative skew)
- $q = 1$: original data
- $q < 1$: reduce concentration on the left (reduce positive skew)
- $\log(x+1)$ may be applied when $x=0$. If distribution of $\log(x+1)$ is normal, it is called lognormal distribution




Histogram of log of income



Boxplot of log of income



Quantile-normal plot of log of income



Source: 2018 American Community Survey.



Skewness and Kurtosis

```
. sum lnincome [fweight=perwt], d
```

		lnincome			
Percentiles		Smallest			
1%	6.214608	1.386294			
5%	7.783224	1.386294			
10%	8.630522	1.386294	Obs	13,849,398	
25%	9.680344	1.386294	Sum of Wgt.	13,849,398	
50%	10.43412		Mean	10.22871	
		Largest		Std. Dev.	1.233225
75%	11.0021	13.05622			
90%	11.51293	13.05622	Variance	1.520844	
95%	11.82041	13.05622	Skewness	-1.123294	
99%	13.05622	13.05622	Kurtosis	5.349345	

Interpretation of $\ln(\text{income})$

(with continuous independent variables)

- With the logarithm of the dependent variable
 - Coefficients are interpreted as percentage changes
- If coefficient of x_1 equals 0.12
 - $\exp(\beta_1)$ times
 - x_1 increases by one unit, y increases on average **1.13 times**, controlling for other independent variables
 - $100 * [\exp(\beta_1) - 1]$ percent
 - x_1 increases by one unit, y increases on average by **13%**, controlling for other independent variables
- If coefficient has a small magnitude: $-0.3 < \beta < 0.3$
 - $100 * \beta$ percent
 - x_1 increases by one unit, y increases on average **approximately by 12%**, controlling for other independents



TEXAS A&M
UNIVERSITY.

$\ln(\text{income}) = F(\text{age}, \text{education})$

```
. ***Use complex survey design
. svy: reg lnincome age educgr
(running regress on estimation sample)
```

Survey: Linear regression

Number of strata	=	212	Number of obs	=	127,785
Number of PSUs	=	79,499	Population size	=	13,849,398
			Design df	=	79,287
			F(2, 79286)	=	7451.80
			Prob > F	=	0.0000
			R-squared	=	0.1932

lnincome	Coef.	Linearized Std. Err.	t	P> t	[95% Conf. Interval]	
age	.0224959	.0003153	71.35	0.000	.0218779	.0231139
educgr	.3381717	.0032453	104.20	0.000	.331811	.3445324
_cons	8.34881	.0175456	475.84	0.000	8.31442	8.383199

Exponential of coefficients

```
. ***Automatically see exponential of coefficients
. svy: reg lnincome age educgr, eform(Exp. Coef.)
(running regress on estimation sample)
```

Survey: Linear regression

Number of strata	=	212	Number of obs	=	127,785
Number of PSUs	=	79,499	Population size	=	13,849,398
			Design df	=	79,287
			F(2, 79286)	=	7451.80
			Prob > F	=	0.0000
			R-squared	=	0.1932

lnincome	Linearized				
	Exp. Coef.	Std. Err.	t	P> t	[95% Conf. Interval]
age	1.022751	.0003225	71.35	0.000	1.022119 1.023383
educgr	1.402381	.0045511	104.20	0.000	1.393489 1.41133
_cons	4225.149	74.13273	475.84	0.000	4082.319 4372.976

Interpretation of age

(income with continuous independent variables)

- Coefficient for **age** equals 0.0225
 - $\exp(\beta_1)$ times
 - When age increases by one unit, income increases on average by **1.0228 times**, controlling for education
 - $100 * [\exp(\beta_1) - 1]$ percent
 - When age increases by one unit, income increases on average by **2.28%**, controlling for education
 - $100 * \beta_1$ percent
 - When age increases by one unit, income increases on average **approximately by 2.25%**, controlling for education



Interpretation of education

(income with continuous independent variables)

- Coefficient for **education** equals 0.3382
 - $\exp(\beta_1)$ times
 - When education increases by one unit, income increases on average by **1.4024 times**, controlling for age
 - $100 * [\exp(\beta_1) - 1]$ percent
 - When education increases by one unit, income increases on average by **40.24%**, controlling for age
 - $100 * \beta_1$ percent
 - When education increases by one unit, income increases on average **approximately by 33.82%**, controlling for age



Standardized coefficients

- . *****Standardized regression coefficients**
- . *****(i.e., standardized partial slopes, beta-weights)**
- . *****It does not allow the use of complex survey design**
- . *****Use pweight to maintain sample size and estimate robust standard errors**
- . **reg lnincome age educgr [pweight=perwt], beta**
(sum of wgt is 13,849,398)

```
Linear regression                Number of obs    =    127,785
                                F(2, 127782)    =    7996.52
                                Prob > F              =    0.0000
                                R-squared              =    0.1932
                                Root MSE           =    1.1077
```

lnincome	Coef.	Robust Std. Err.	t	P> t	Beta
age	.0224959	.0002969	75.76	0.000	.2637902
educgr	.3381717	.0031694	106.70	0.000	.3246429
_cons	8.34881	.0166508	501.41	0.000	.

Interpretation of standardized

(income with continuous independent variables)

- Coefficient for **age** equals 0.2638
 - $\exp(\beta_1)$ times
 - When age increases by one standard deviation, income increases on average by **1.3019 times**, controlling for education
 - $100 * [\exp(\beta_1) - 1]$ percent
 - When age increases by one standard deviation, income increases on average by **30.19%**, controlling for education
 - $100 * \beta_1$ percent
 - When age increases by one standard deviation, income increases on average **approximately by 26.38%**, controlling for education

Adjusted R^2

```
. ***Use aweight to estimate adjusted R-squared
. ***pweight and complex survey design omit sum of squares and adjusted R-squared
. reg lnincome age educgr [aweight=perwt]
(sum of wgt is 13,849,398)
```

Source	SS	df	MS	Number of obs	=	127,785
Model	37544.8387	2	18772.4194	F(2, 127782)	=	15298.69
Residual	156796.221	127,782	1.22706031	Prob > F	=	0.0000
Total	194341.059	127,784	1.52085597	R-squared	=	0.1932
				Adj R-squared	=	0.1932
				Root MSE	=	1.1077

lnincome	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
age	.0224959	.0002155	104.39	0.000	.0220735	.0229183
educgr	.3381717	.0026324	128.47	0.000	.3330122	.3433311
_cons	8.34881	.0113381	736.35	0.000	8.326587	8.371032



TEXAS A&M
UNIVERSITY.

Predicted values

- We can estimate the predicted values of the dependent variable for each individual in the dataset
- Use the estimated coefficients from the regression model

$$y_i' = \hat{y}_i = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i}$$

Predicted income

- Income = F(age, education)

income	Linearized			P> t	[95% Conf. Interval]	
	Coef.	Std. Err.	t			
age	796.3443	11.73077	67.89	0.000	773.3521	819.3366
educgr	16863.33	179.705	93.84	0.000	16511.11	17215.55
_cons	-31880.99	661.937	-48.16	0.000	-33178.38	-30583.59

- Use the regression equation to predict income for someone with 45 years of age and college education

$$\hat{y} = -31,880.99 + 796.34(\text{age}) + 16,863.33(\text{educgr})$$

$$\hat{y} = -31,880.99 + (796.34)(45) + (16,863.33)(4)$$

$$\hat{y} = 71,407.63$$

- Under these conditions, we would predict 71,407.63 dollars for that individual




Microdata


	age	educgr	income	predincome
1	21	2	3200	18568.9
2	20	2	35000	17772.56
3	31	2	10000	26532.34
4	39	4	30000	66629.76
5	18	2	1500	16179.87
6	25	1	13000	4890.951
7	20	3	5600	34635.88
8	34	2	65000	28921.38
9	18	2	4000	16179.87
10	18	3	1400	33043.2
11	20	2	5000	17772.56
12	18	2	2300	16179.87
13	20	2	18000	17772.56
14	19	3	14000	33839.54
15	20	2	6000	17772.56
16	19	2	1800	16976.21
17	21	3	320	35432.23
18	22	3	1900	36228.57
19	46	2	28000	38477.51
20	20	3	5000	34635.88
21	23	3	1000	37024.92
22	19	2	10000	16976.21
23	19	3	600	33839.54
24	20	3	10000	34635.88
25	22	3	7000	36228.57
26	22	3	4000	36228.57
27	48	3	11000	56933.53
28	23	3	140	37024.92
29	21	3	2000	35432.23
30	21	3	3600	35432.23



Predicted income by age



Predicted income by education



Predicted log of income

- $\ln(\text{income}) = F(\text{age}, \text{education})$

lnincome	Linearized					[95% Conf. Interval]	
	Coef.	Std. Err.	t	P> t			
age	.0224959	.0003153	71.35	0.000	.0218779	.0231139	
educgr	.3381717	.0032453	104.20	0.000	.331811	.3445324	
_cons	8.34881	.0175456	475.84	0.000	8.31442	8.383199	

- Use the regression equation to predict log of income for someone with 45 years of age and college education

$$\ln(\hat{y}) = 8.3488 + 0.0225(\text{age}) + 0.3382(\text{educgr})$$

$$\ln(\hat{y}) = 8.3488 + (0.0225)(45) + (0.3382)(4)$$

$$\ln(\hat{y}) = 10.7141$$

$$\hat{y} = 44,985.70$$

- Under these conditions, we would predict 44,985.70 dollars for that individual




Microdata


	age	educgr	income	lnincome	predlnincome
1	21	2	3200	8.070906	9.497567
2	20	2	35000	10.4631	9.475071
3	31	2	10000	9.21034	9.722527
4	39	4	30000	10.30895	10.57884
5	18	2	1500	7.313221	9.430079
6	25	1	13000	9.472705	9.249379
7	20	3	5600	8.630522	9.813243
8	34	2	65000	11.08214	9.790014
9	18	2	4000	8.294049	9.430079
10	18	3	1400	7.244227	9.768251
11	20	2	5000	8.517193	9.475071
12	18	2	2300	7.740664	9.430079
13	20	2	18000	9.798127	9.475071
14	19	3	14000	9.546813	9.790747
15	20	2	6000	8.699514	9.475071
16	19	2	1800	7.495542	9.452576
17	21	3	320	5.768321	9.835739
18	22	3	1900	7.549609	9.858234
19	46	2	28000	10.23996	10.05997
20	20	3	5000	8.517193	9.813243
21	23	3	1000	6.907755	9.880731
22	19	2	10000	9.21034	9.452576
23	19	3	600	6.39693	9.790747
24	20	3	10000	9.21034	9.813243
25	22	3	7000	8.853665	9.858234
26	22	3	4000	8.294049	9.858234
27	48	3	11000	9.305651	10.44313
28	23	3	140	4.941642	9.880731
29	21	3	2000	7.600903	9.835739
30	21	3	3600	8.188689	9.835739




Predicted $\ln(\text{income})$ by age




Exponential of predicted $\ln(\text{income})$ by age




Predicted $\ln(\text{income})$ by education



Exponential of predicted $\ln(\text{income})$ by education





TEXAS A&M
UNIVERSITY.

Residual analysis with graphs

- Homoscedasticity assumption
 - The variance of y scores is uniform for all values of x
 - If the y scores are evenly spread above and below the regression line for the entire length of the line, the association is homoscedastic
- The same assumption applies to residuals
 - Difference between observed value (y) and predicted value (\hat{y})
 - $e = y - \hat{y}$
 - We can plot residuals against predicted values \hat{y} (which summarize all x variables)



Microdata

	age	educgr	income	predincome	resincome	lnincome	predlnincome	reslnincome
1	21	2	3200	18568.9	-15368.9	8.070906	9.497567	-1.426661
2	20	2	35000	17772.56	17227.44	10.4631	9.475071	.9880321
3	31	2	10000	26532.34	-16532.34	9.21034	9.722527	-.5121856
4	39	4	30000	66629.76	-36629.75	10.30895	10.57884	-.2698844
5	18	2	1500	16179.87	-14679.87	7.313221	9.430079	-2.116859
6	25	1	13000	4890.951	8109.049	9.472705	9.249379	.2233258
7	20	3	5600	34635.88	-29035.88	8.630522	9.813243	-1.182721
8	34	2	65000	28921.38	36078.62	11.08214	9.790014	1.292129
9	18	2	4000	16179.87	-12179.87	8.294049	9.430079	-1.13603
10	18	3	1400	33043.2	-31643.2	7.244227	9.768251	-2.524024
11	20	2	5000	17772.56	-12772.56	8.517193	9.475071	-.9578784
12	18	2	2300	16179.87	-13879.87	7.740664	9.430079	-1.689415
13	20	2	18000	17772.56	227.4432	9.798127	9.475071	.323056
14	19	3	14000	33839.54	-19839.54	9.546813	9.790747	-.243934
15	20	2	6000	17772.56	-11772.56	8.699514	9.475071	-.7755568
16	19	2	1800	16976.21	-15176.21	7.495542	9.452576	-1.957033
17	21	3	320	35432.23	-35112.23	5.768321	9.835739	-4.067418
18	22	3	1900	36228.57	-34328.57	7.549609	9.858234	-2.308625
19	46	2	28000	38477.51	-10477.51	10.23996	10.05997	.179995
20	20	3	5000	34635.88	-29635.88	8.517193	9.813243	-1.29605
21	23	3	1000	37024.92	-36024.92	6.907755	9.880731	-2.972975
22	19	2	10000	16976.21	-6976.212	9.21034	9.452576	-.2422348
23	19	3	600	33839.54	-33239.54	6.39693	9.790747	-3.393817
24	20	3	10000	34635.88	-24635.88	9.21034	9.813243	-.6029024
25	22	3	7000	36228.57	-29228.57	8.853665	9.858234	-1.004569
26	22	3	4000	36228.57	-32228.57	8.294049	9.858234	-1.564185
27	48	3	11000	56933.53	-45933.53	9.305651	10.44313	-1.137478
28	23	3	140	37024.92	-36884.92	4.941642	9.880731	-4.939088
29	21	3	2000	35432.23	-33432.23	7.600903	9.835739	-2.234836
30	21	3	3600	35432.23	-31832.23	8.188689	9.835739	-1.64705









Figure 2.10 "All clear" e -versus- \hat{Y} plot (artificial data).




Influential Case



Curvilinear Relation




Nonnormal Residual Distribution




Heteroscedasticity

Figure 2.11 Examples of trouble seen in e -versus- \hat{Y} plots (artificial data).

Residuals: $\text{Income} = F(\text{age, education})$



Residuals: $\ln(\text{income}) = F(\text{age}, \text{education})$





TEXAS A&M
UNIVERSITY.

OLS with age and age squared

- $\ln(\text{income})$ as a function of age and age squared

$$y = \beta_0 + \beta_1 x + \beta_2 x^2 + u$$

- Variation in income due to variation in age


$$\Delta y / \Delta x \approx \beta_1 + 2\beta_2 x$$

- Marginal effect of age on income depends on β_1 , β_2 , and specific age value (x)
- There is a positive value of x , in which the effect of x on y is zero, called the critical point (x^*)

$$x^* = |\beta_1 / (2\beta_2)|$$



Mean income by age



Source: 2018 American Community Survey.



$\ln(\text{income}) = F(\text{age}, \text{age squared})$

```
. ***OLS with natural logarithm of income, age, and age squared
. svy: reg lnincome age agesq
(running regress on estimation sample)
```

Survey: Linear regression

Number of strata	=	212	Number of obs	=	127,785
Number of PSUs	=	79,499	Population size	=	13,849,398
			Design df	=	79,287
			F(2, 79286)	=	7983.37
			Prob > F	=	0.0000
			R-squared	=	0.2185

lnincome	Coef.	Linearized Std. Err.	t	P> t	[95% Conf. Interval]	
age	.1943162	.0017962	108.18	0.000	.1907956	.1978369
agesq	-.0019721	.0000205	-96.06	0.000	-.0020123	-.0019319
_cons	6.009389	.0368055	163.27	0.000	5.937251	6.081528

Association of income with age

- Variation in income due to variation in age

$$\Delta \ln(\text{income}) / \Delta \text{age} \approx \beta_1 + 2\beta_2(\text{age})$$

$$\Delta \ln(\text{income}) / \Delta \text{age} \approx 0.1943 + 2(-0.0020)(\text{age})$$

$$\Delta \ln(\text{income}) / \Delta \text{age} \approx 0.1943 - 0.0040(\text{age})$$


- Critical point (curve changes from upward to downward)

$$\text{age}^* = |\beta_1 / (2\beta_2)| = |0.1943 / (2 \cdot -0.0020)|$$


$$\text{age}^* = |-48.57| = 48.57$$




Predicted $\ln(\text{income})$ by age, age^2




Exponential of predicted $\ln(\text{income})$ by age, age^2




Residuals: $\ln(\text{income}) = F(\text{age}, \text{age}^2)$



Residuals: $\text{Exp. } \ln(\text{income}) = F(\text{age}, \text{age}^2)$





TEXAS A&M
UNIVERSITY.

Dummy variables


- Many variables that are important in social life are nominal-level variables
 - They cannot be included in a regression equation or correlational analysis (e.g., sex, race/ethnicity)
- We can create dummy variables
 - Two categories, one coded as 0 and the other as 1

Sex	Male	Female
1	1	0
2	0	1

Race/ ethnicity	White	Black	Hispanic	Other
1	1	0	0	0
2	0	1	0	0
3	0	0	1	0
4	0	0	0	1

Age in interval-ratio level

- Age does not have a normal distribution



- Generate age group variable (categorical)
 - 16–19; 20–24; 25–34; 35–44; 45–54; 55–64; 65+



Age in ordinal level

- Age has seven categories

```
. table agegr, contents(min age max age count age)
```

agegr	min(age)	max(age)	N(age)
16	16	19	6,337
20	20	24	11,945
25	25	34	26,752
35	35	44	25,575
45	45	54	25,454
55	55	64	22,457
65	65	92	9,265

- Generate dummy variables for age...



Dummies for age

- Generate dummy variables for age group

Age group	Age 16–19	Age 20–24	Age 25–34	Age 35–44	Age 45–54	Age 55–64	Age 65+
16–19	1	0	0	0	0	0	0
20–24	0	1	0	0	0	0	0
25–34	0	0	1	0	0	0	0
35–44	0	0	0	1	0	0	0
45–54	0	0	0	0	1	0	0
55–64	0	0	0	0	0	1	0
65+	0	0	0	0	0	0	1

Reference category

- Use the category with the largest sample size as the reference (25–34)

`. tab agegr, m`

agegr	Freq.	Percent	Cum.
16	6,337	4.96	4.96
20	11,945	9.35	14.31
25	26,752	20.94	35.24
35	25,575	20.01	55.26
45	25,454	19.92	75.18
55	22,457	17.57	92.75
65	9,265	7.25	100.00
Total	127,785	100.00	

- Or category with large sample and meaningful interpretation for your problem (age group with the highest average income: 45–54)


`. table agegr, c(mean income)`

agegr	mean(income)
16	6051.891
20	18397.36
25	42752.68
35	61426.85
45	67367.77
55	65728.8
65	50250.71



Educational attainment

- Education does not have a normal distribution



- Generate education group variable (categorical)
 - Less than high school; high school; some college; college; graduate school



Education in ordinal level

- Education has five categories

```
. tab educgr, m
```

educgr	Freq.	Percent	Cum.
Less than high school	12,719	9.95	9.95
High school	40,869	31.98	41.94
Some college	30,360	23.76	65.69
College	28,110	22.00	87.69
Graduate school	15,727	12.31	100.00
Total	127,785	100.00	

- Generate dummy variables for education...



Dummies for education

- Generate dummy variables for education group

Education group	<High school	High school	Some College	College	Graduate school
Less than high school	1	0	0	0	0
High school	0	1	0	0	0
Some college	0	0	1	0	0
College	0	0	0	1	0
Graduate school	0	0	0	0	1



Reference group

- Use the category with the largest sample size as the reference (high school)

```
. tab educgr, m
```

educgr	Freq.	Percent	Cum.
Less than high school	12,719	9.95	9.95
High school	40,869	31.98	41.94
Some college	30,360	23.76	65.69
College	28,110	22.00	87.69
Graduate school	15,727	12.31	100.00
Total	127,785	100.00	



log income = F(age, education)

```
. svy: reg lnincome ib45.agegr ib2.educgr
(running regress on estimation sample)
```

Survey: Linear regression

Number of strata	=	212	Number of obs	=	127,785
Number of PSUs	=	79,499	Population size	=	13,849,398
			Design df	=	79,287
			F(10, 79278)	=	2860.65
			Prob > F	=	0.0000
			R-squared	=	0.3129

lnincome	Linearized					[95% Conf. Interval]
	Coef.	Std. Err.	t	P> t		
agegr						
16	-2.223012	.0227431	-97.74	0.000	-2.267588	-2.178435
20	-1.151434	.0155642	-73.98	0.000	-1.18194	-1.120928
25	-.3856507	.0104177	-37.02	0.000	-.4060693	-.365232
35	-.0929935	.0104004	-8.94	0.000	-.1133781	-.0726089
55	-.053233	.0111394	-4.78	0.000	-.0750662	-.0313998
65	-.5928305	.0186409	-31.80	0.000	-.6293667	-.5562944
educgr						
Less than high school	-.3066773	.0128821	-23.81	0.000	-.3319261	-.2814286
Some college	.1354166	.0097974	13.82	0.000	.1162138	.1546194
College	.5445375	.0101702	53.54	0.000	.524604	.564471
Graduate school	.8187744	.0121	67.67	0.000	.7950584	.8424904
_cons	10.41295	.0092523	1125.44	0.000	10.39482	10.43109

Exponential of coefficients

```
. ***Automatically see exponential of coefficients
. svy: reg lnincome ib45.agegr ib2.educgr, eform(Exp. Coef.)
(running regress on estimation sample)
```

Survey: Linear regression

Number of strata	=	212	Number of obs	=	127,785
Number of PSUs	=	79,499	Population size	=	13,849,398
			Design df	=	79,287
			F(10, 79278)	=	2860.65
			Prob > F	=	0.0000
			R-squared	=	0.3129

lnincome	Linearized					[95% Conf. Interval]
	Exp. Coef.	Std. Err.	t	P> t		
agegr						
16	.1082825	.0024627	-97.74	0.000	.1035617	.1132186
20	.316183	.0049211	-73.98	0.000	.3066833	.325977
25	.680008	.0070841	-37.02	0.000	.666264	.6940356
35	.9111994	.0094768	-8.94	0.000	.892813	.9299645
55	.948159	.0105619	-4.78	0.000	.9276821	.969088
65	.5527605	.010304	-31.80	0.000	.5329292	.5733297
educgr						
Less than high school	.735888	.0094797	-23.81	0.000	.7175404	.7547048
Some college	1.145014	.0112182	13.82	0.000	1.123236	1.167214
College	1.723811	.0175315	53.54	0.000	1.68979	1.758517
Graduate school	2.267719	.0274395	67.67	0.000	2.21457	2.322143
_cons	33288.07	307.9918	1125.44	0.000	32689.85	33897.24



Interpretation of age

(log of income with dummies as independent variables)

- 45–54 age group is reference category for **age**
- Coefficient for 16–19 age group equals -2.2230
 - $\exp(\beta_1)$ times
 - People between 16–19 years of age have on average earnings **0.1083 times** the earnings of people between 45–54 years of age, controlling for the other independent variables
 - $100 * [\exp(\beta_1) - 1]$ percent
 - People between 16–19 years of age have on average earnings **89.17% lower** than earnings of people between 45–54 years of age, controlling for the other independent variables
 - $100 * \beta_1$ percent: result is not good because $\beta_1 > 0.3$
 - People between 16–19 years of age have on average earnings **approximately 222.30% lower** than earnings of people between 45–54 years of age, controlling for the other independent variables

Interpretation of education

(log of income with dummies as independent variables)

- High school is reference category for **education**
- Coefficient for college equals 0.5445
 - $\exp(\beta_1)$ times
 - People with college degree have on average earnings **1.7238 times higher** than earnings of high school graduates, controlling for the other independent variables
 - $100 * [\exp(\beta_1) - 1]$ percent
 - People with college degree have on average earnings **72.38% higher** than earnings of high school graduates, controlling for the other independent variables
 - $100 * \beta_1$ percent: result is not good because $\beta_1 > 0.3$
 - People with college degree have on average earnings **approximately 54.45% higher** than earnings of high school graduates, controlling for the other independent variables

Standardized coefficients

```
. reg lnincome ib45.agegr ib2.educgr [pweight=perwt], beta
(sum of wgt is 13,849,398)
```


Linear regression

```
Number of obs   =   127,785
F(10, 127774)   =   3037.91
Prob > F        =   0.0000
R-squared       =   0.3129
Root MSE       =   1.0223
```


lnincome	Coef.	Robust Std. Err.	t	P> t	Beta
agegr					
16	-2.223012	.022166	-100.29	0.000	-.3875416
20	-1.151434	.0148555	-77.51	0.000	-.290206
25	-.3856507	.0103423	-37.29	0.000	-.1333188
35	-.0929935	.0103849	-8.95	0.000	-.0310561
55	-.053233	.0110966	-4.80	0.000	-.0151658
65	-.5928305	.018443	-32.14	0.000	-.107231
educgr					
Less than high school	-.3066773	.0125263	-24.48	0.000	-.0788327
Some college	.1354166	.0096013	14.10	0.000	.047455
College	.5445375	.0100048	54.43	0.000	.1781623
Graduate school	.8187744	.0120082	68.18	0.000	.2068187
_cons	10.41295	.0091286	1140.69	0.000	.




Residuals: $\ln(\text{income}) = F(\text{age group}, \text{educ. group})$



Residuals: $\text{Exp. } \ln(\text{income}) = F(\text{age group, educ. group})$





TEXAS A&M
UNIVERSITY.

Full OLS model

- Dependent variable
 - Natural logarithm of income
- Independent variables
 - **Sex:** female; male (reference)
 - **Age group:** 16–19; 20–24; 25–34; 35–44; 45–54 (reference); 55–64; 65+
 - **Education group:** less than high school, high school (reference), some college, college, graduate school
 - **Race/ethnicity:** White (reference); African American; Hispanic; Asian; Native American; Other races
 - **Marital status:** married (reference); separated, divorced, widowed; never married
 - **Migration status:** non-migrant (reference); internal migrant; international migrant

Command in Stata

```
. svy: reg lnincome i.female ib45.agegr ib2.educgr i.raceth i.marital i.migrant  
(running regress on estimation sample)
```

Survey: Linear regression

Number of strata	=	212	Number of obs	=	127,785
Number of PSUs	=	79,499	Population size	=	13,849,398
			Design df	=	79,287
			F(20, 79268)	=	1818.83
			Prob > F	=	0.0000
			R-squared	=	0.3577



Coefficients
from OLS
regression for
natural logarithm
of income,
Texas, 2018

lnincome	Coef.	Linearized Std. Err.	t	P> t	[95% Conf. Interval]	
female						
Female	-.4374635	.0070675	-61.90	0.000	-.4513158	-.4236111
agegr						
16-19	-1.995369	.0241877	-82.50	0.000	-2.042777	-1.947961
20-24	-.9592868	.0168846	-56.81	0.000	-.9923806	-.926193
25-34	-.2920554	.0106538	-27.41	0.000	-.3129368	-.271174
35-44	-.0705981	.0100164	-7.05	0.000	-.0902301	-.0509661
55-64	-.0751899	.0107209	-7.01	0.000	-.0962027	-.0541771
65-100	-.6377643	.0183047	-34.84	0.000	-.6736413	-.6018873
educgr						
Less than high school	-.3148089	.01281	-24.58	0.000	-.3399165	-.2897013
Some college	.1565395	.0096239	16.27	0.000	.1376767	.1754023
College	.5426535	.0101186	53.63	0.000	.5228211	.562486
Graduate school	.8081078	.0122256	66.10	0.000	.7841457	.8320698
raceth						
African American	-.172703	.012575	-13.73	0.000	-.19735	-.148056
Hispanic	-.1285316	.0085376	-15.05	0.000	-.1452652	-.111798
Asian	-.1583612	.0172829	-9.16	0.000	-.1922356	-.1244867
Native American	-.071535	.0555021	-1.29	0.197	-.1803187	.0372488
Other races	-.1193284	.0302909	-3.94	0.000	-.1786982	-.0599585
marital						
Separated, divorced, wid..	-.1364001	.0101838	-13.39	0.000	-.1563603	-.11644
Never married	-.2696217	.009485	-28.43	0.000	-.2882122	-.2510312
migrant						
Internal migrant	-.1211724	.0160131	-7.57	0.000	-.1525579	-.0897869
International migrant	-.4936644	.0683904	-7.22	0.000	-.6277092	-.3596197
_cons	10.76426	.0105691	1018.47	0.000	10.74355	10.78498

Exponential of coefficients from OLS regression for natural logarithm of income, Texas, 2018

lnincome	Exp. Coef.	Linearized Std. Err.	t	P> t	[95% Conf. Interval]	
female						
Female	.6456721	.0045633	-61.90	0.000	.6367897	.6546784
agegr						
16-19	.1359635	.0032886	-82.50	0.000	.1296682	.1425645
20-24	.3831661	.0064696	-56.81	0.000	.3706932	.3960586
25-34	.7467272	.0079555	-27.41	0.000	.7312961	.7624838
35-44	.9318363	.0093336	-7.05	0.000	.9137209	.9503108
55-64	.9275673	.0099443	-7.01	0.000	.9082799	.9472643
65-100	.5284726	.0096735	-34.84	0.000	.5098487	.5477769
educgr						
Less than high school	.7299283	.0093504	-24.58	0.000	.7118298	.7484871
Some college	1.169457	.0112548	16.27	0.000	1.147604	1.191726
College	1.720566	.0174098	53.63	0.000	1.686779	1.75503
Graduate school	2.243658	.02743	66.10	0.000	2.190535	2.29807
raceth						
African American	.8413875	.0105805	-13.73	0.000	.8209033	.8623828
Hispanic	.8793858	.0075078	-15.05	0.000	.8647929	.8942249
Asian	.8535415	.0147517	-9.16	0.000	.8251124	.88295
Native American	.9309637	.0516704	-1.29	0.197	.835004	1.037951
Other races	.8875163	.0268836	-3.94	0.000	.8363582	.9418036
marital						
Separated, divorced, widowed	.8724934	.0088853	-13.39	0.000	.855251	.8900835
Never married	.7636683	.0072434	-28.43	0.000	.7496025	.7779981
migrant						
Internal migrant	.8858812	.0141857	-7.57	0.000	.8585092	.9141259
International migrant	.6103856	.0417445	-7.22	0.000	.5338133	.6979417
_cons	47299.9	499.9155	1018.47	0.000	46330.15	48289.95

Edited table

Table 1. Coefficients and standard errors estimated with ordinary least squares models for the logarithm of wage and salary income as the dependent variable, Texas, 2018

Independent variables	Model 1	Model 2	Model 3	Model 4	Model 4 Standardized coefficients
Constant	10.61*** (0.00961)	10.70*** (0.0106)	10.76*** (0.0106)	10.76*** (0.0106)	
Sex					
Male	ref.	ref.	ref.	ref.	ref.
Female	-0.449*** (0.00700)	-0.444*** (0.00700)	-0.436*** (0.00707)	-0.437*** (0.00707)	-0.177
Age groups					
16-19	-2.195*** (0.0226)	-2.204*** (0.0228)	-2.007*** (0.0241)	-1.995*** (0.0242)	-0.348
20-24	-1.154*** (0.0155)	-1.142*** (0.0155)	-0.973*** (0.0168)	-0.959*** (0.0169)	-0.242
25-34	-0.396*** (0.0103)	-0.385*** (0.0102)	-0.302*** (0.0106)	-0.292*** (0.0107)	-0.101
35-44	-0.100*** (0.0101)	-0.0921*** (0.0101)	-0.0734*** (0.0100)	-0.0706*** (0.0100)	-0.0236
45-54	ref.	ref.	ref.	ref.	ref.
55-64	-0.0545*** (0.0108)	-0.0698*** (0.0108)	-0.0737*** (0.0107)	-0.0752*** (0.0107)	-0.0214
65+	-0.604*** (0.0183)	-0.631*** (0.0183)	-0.634*** (0.0183)	-0.638*** (0.0183)	-0.115

Note: Coefficients and standard errors were generated with the complex survey design of the American Community Survey. The standardized coefficients were generated with sample weights. Standard errors are reported in parentheses. *Significant at p<0.10; **Significant at p<0.05; ***Significant at p<0.01.

Source: 2018 American Community Survey.

Continue...

Edited table

Table 1. Coefficients and standard errors estimated with ordinary least squares models for the logarithm of wage and salary income as the dependent variable, Texas, 2018

Independent variables	Model 1	Model 2	Model 3	Model 4	Model 4 Standardized coefficients
Education groups					
Less than high school	-0.336*** (0.0125)	-0.311*** (0.0129)	-0.314*** (0.0128)	-0.315*** (0.0128)	-0.0809
High school	ref.	ref.	ref.	ref.	ref.
Some college	0.165*** (0.00965)	0.156*** (0.00971)	0.157*** (0.00963)	0.157*** (0.00962)	0.0549
College	0.579*** (0.0100)	0.551*** (0.0102)	0.539*** (0.0101)	0.543*** (0.0101)	0.178
Graduate school	0.848*** (0.0119)	0.826*** (0.0123)	0.803*** (0.0122)	0.808*** (0.0122)	0.204

Note: Coefficients and standard errors were generated with the complex survey design of the American Community Survey. The standardized coefficients were generated with sample weights. Standard errors are reported in parentheses. *Significant at p<0.10; **Significant at p<0.05; ***Significant at p<0.01.

Source: 2018 American Community Survey.

Edited table

Table 1. Coefficients and standard errors estimated with ordinary least squares models for the logarithm of wage and salary income as the dependent variable, Texas, 2018

Independent variables	Model 1	Model 2	Model 3	Model 4	Model 4 Standardized coefficients
Race/ethnicity					
White		ref.	ref.	ref.	ref.
African American		-0.211*** (0.0126)	-0.172*** (0.0126)	-0.173*** (0.0126)	-0.0461
Hispanic		-0.132*** (0.00860)	-0.125*** (0.00853)	-0.129*** (0.00854)	-0.0503
Asian		-0.153*** (0.0176)	-0.166*** (0.0175)	-0.158*** (0.0173)	-0.0288
Native American		-0.0988* (0.0540)	-0.0758 (0.0549)	-0.0715 (0.0555)	-0.00272
Other races		-0.140*** (0.0302)	-0.124*** (0.0301)	-0.119*** (0.0303)	-0.0123

Note: Coefficients and standard errors were generated with the complex survey design of the American Community Survey. The standardized coefficients were generated with sample weights. Standard errors are reported in parentheses. *Significant at $p < 0.10$; **Significant at $p < 0.05$; ***Significant at $p < 0.01$.

Source: 2018 American Community Survey.

Edited table

Table 1. Coefficients and standard errors estimated with ordinary least squares models for the logarithm of wage and salary income as the dependent variable, Texas, 2018


Independent variables	Model 1	Model 2	Model 3	Model 4	Model 4 Standardized coefficients
Marital status					
Married			ref.	ref.	ref.
Separated, divorced, widowed			-0.139*** (0.0102)	-0.136*** (0.0102)	-0.0398
Never married			-0.270*** (0.00950)	-0.270*** (0.00948)	-0.104
Migration status					
Non-migrant				ref.	ref.
Internal migrant				-0.121*** (0.0160)	-0.0242
International migrant				-0.494*** (0.0684)	-0.0287
R ²	0.346	0.349	0.356	0.358	0.358
Observations	127,785	127,785	127,785	127,785	127,785

Note: Coefficients and standard errors were generated with the complex survey design of the American Community Survey. The standardized coefficients were generated with sample weights. Standard errors are reported in parentheses. *Significant at p<0.10; **Significant at p<0.05; ***Significant at p<0.01.

Source: 2018 American Community Survey.


Exponential of age group coefficients

(Example of how to show regression results in conferences. Edited in Excel)



Exponential of age group coefficients


(Example of how to show regression results in conferences. Edited in Excel.)



Predicted female income by age

(Using “mgen” command within SPost13 package by Long and Freese, 2014)


```
mgen, stub(F) at(agegr=(16 20 25 35 45 55 65) female=1 ///  
    raceth=1 educgr=2 marital=1 migrant=1) allstats
```



Predicted male income by age


(Using “mgen” command within SPost13 package by Long and Freese, 2014)

```
mgen, stub(M) at(agegr=(16 20 25 35 45 55 65) female=0 ///  
    raceth=1 educgr=2 marital=1 migrant=1) allstats
```




Predicted income by age and sex

For White, High School, Married, Non-migrant




Predicted income by age and sex

For White, High School, Married, Non-migrant




Residuals


$$\ln(\text{income}) = F(\text{sex}, \text{age}, \text{educ}, \text{race/ethnicity}, \text{marital}, \text{migrant})$$



Residuals

$\text{Exp.} \ln(\text{income}) = F(\text{sex, age, educ, race/ethnicity, marital, migrant})$






TEXAS A&M
UNIVERSITY.

Interaction with dummy variables

- As before, we can simply include dummy variables as independent variables

$$\text{earnings} = \beta_0 + \delta_0 \text{ women} + \beta_1 \text{ education} + u$$

- Difference between sexes does not depend on the level of education (fitted lines are parallel)




Different slopes


- We can test if the effect of education on earnings vary by sex


$$\text{earnings} = (\beta_0 + \delta_0 \text{ women}) + (\beta_1 + \delta_1 \text{ women}) * \text{educ} + u$$

$$\delta_0 < 0, \delta_1 < 0$$



$$\delta_0 < 0, \delta_1 > 0$$





TEXAS A&M
UNIVERSITY.

Age-education & earnings, Brazil


Year	Area	Log of mean earnings	Age-education group	Dummies for age-education groups			Distr. of male pop.	Variables for distribution of male population			Num. of obs.
		log(Y_{git})	G11–G44	G11	...	G44	P11–P44	P11	...	P44	
1970	110006	5.80	15–24 years & < primary	1	...	0	0.221	0.221	...	0	2,016
1970	110006	6.02	15–24 years & primary	0	...	0	0.102	0	...	0	927
1970	110006	6.57	15–24 years & secondary	0	...	0	0.007	0	...	0	62
1970	110006	7.58	15–24 years & university	0	...	0	0.001	0	...	0	11
...
1970	110006	7.91	50–64 years & university	0	...	1	0.002	0.002	15
...

Fixed effects models


	Baseline model	Composition model
Dependent variable		
Logarithm of the mean real monthly earnings by age-education group, area, and time	$\log(Y_{git})$	$\log(Y_{git})$
Independent variables		
16 age-education indicators * time	$(G_{11}-G_{44}) * \theta_t$	$(G_{11}-G_{44}) * \theta_t$
Distribution of male population into 16 age-education groups * time		$(P_{11}-P_{44}) * \theta_t$
Area-time fixed effects	α_{it}	α_{it}

Effects of age-education indicators ($G_{11}-G_{44}$)

Baseline model, Brazil, 2010




Effects of age-education indicators ($G_{11}-G_{44}$) Composition model, Brazil, 2010



Source: 1970, 1980, 1991, 2000, and 2010 Brazilian Demographic Censuses.


Effects of group proportions ($P_{11}-P_{14}$) on earnings, Brazil, 1970–2010

15–24 years




Effects of group proportions ($P_{21}-P_{24}$) on earnings, Brazil, 1970–2010

25–34 years





Effects of group proportions ($P_{31}-P_{34}$) on earnings, Brazil, 1970–2010

35–49 years



Effects of group proportions (P_{41} – P_{44}) on earnings, Brazil, 1970–2010 50–64 years





TEXAS A&M
UNIVERSITY.

Religion & earnings, Brazil

- Unit of analysis
 - Group defined by age, education, area, year
($4 \times 3 \times 502 \times 4 = 24,096$)
- Dependent variable
 - Logarithm of average earnings of each group
- Independent variables
 - Age-education indicators
 - Proportion of Protestants in each group * year
 - Area and year fixed effects




Earnings by proportion Protestants

Prop. Protestants


15–24 years, 0–4 education

○ 1970 x 2000




Prop. Protestants * Year

15–24 years, 0–4 education




Prop. Protestants

25–34 years, 0–4 education



Prop. Protestants * Year

25–34 years, 0–4 education




Earnings by proportion Protestants

Prop. Protestants


35–49 years, 0–4 education

○ 1970 x 2000




Prop. Protestants * Year

35–49 years, 0–4 education




Prop. Protestants

50–64 years, 0–4 education



Prop. Protestants * Year

50–64 years, 0–4 education



Interaction of religion and race


Table 4. Area and Time Fixed-Effects Estimates of Equation With Age-Education Group Indicators, Proportion Protestant, Proportion of Non-Whites, Age-Education Group Indicators Interacted with Year and Region, and Proportion of Protestants Interacted with Proportion of Non-Whites, 1980–2000. Dependent Variable Is $\log(\text{monthly earnings})^\dagger$

Coefficients [‡]	Proportion Protestant	Proportion of non-whites	Protestant *non-white
Ages 15–24 years; 0–4 years of schooling	–0.035 (0.2492)	–0.787*** (0.0581)	0.918 (0.4784)
Ages 25–34 years; 0–4 years of schooling	–0.003 (0.2174)	–0.879*** (0.0575)	1.041* (0.4369)
Ages 35–49 years; 0–4 years of schooling	–0.011 (0.1986)	–0.950*** (0.0583)	1.463** (0.4230)
Ages 50–64 years; 0–4 years of schooling	–0.158 (0.1757)	–0.967*** (0.0565)	1.528*** (0.3765)




Predicted relative earnings of males with 0–4 years of schooling by non-white and Protestant proportions, Brazil, 1980–2000

15–24 years




—■— 0.0 Non-White —◆— 0.3 Non-White —●— 0.6 Non-White —▲— 0.9 Non-White

25–34 years




—■— 0.0 Non-White —◆— 0.3 Non-White —●— 0.6 Non-White —▲— 0.9 Non-White

35–49 years




—■— 0.0 Non-White —◆— 0.3 Non-White —●— 0.6 Non-White —▲— 0.9 Non-White

50–64 years



—■— 0.0 Non-White —◆— 0.3 Non-White —●— 0.6 Non-White —▲— 0.9 Non-White



TEXAS A&M
UNIVERSITY.