Lecture 11: Chi square

Ernesto F. L. Amaral

March 28, 2023
Advanced Methods of Social Research (SOCI 420)

www.ernestoamaral.com

Source: Healey, Joseph F. 2015. "Statistics: A Tool for Social Research." Stamford: Cengage Learning. 10th edition. Chapter 11 (pp. 276–306).



Outline

- Identify and cite examples of situations in which the chi square test is appropriate
- Explain the structure of a bivariate table and the concept of independence as applied to expected and observed frequencies in a bivariate table
- Explain the logic of hypothesis testing in terms of chi square
- Perform the chi square test using the five-step model and correctly interpret the results
- Explain the limitations of the chi square test and, especially, the difference between statistical significance and substantive significance (importance, magnitude)

The bivariate table

 Bivariate tables display the scores of cases on two different variables at the same time

Rates of Participation in Voluntary Associations by Marital Status for 100 Senior Citizens

| | Marita | | | |
|---------------------|---------|-----------|---|------------------|
| Participation Rates | Married | Unmarried | | TOTALS |
| High | | | N | 50 |
| Low | | | | <u>50</u> 100 |
| TOTALS | 50 | 50 | | 100 |



Aspects of the table

- Note the two dimensions: rows and columns
- What is the independent variable?
- What is the dependent variable?
- Where are the row and column marginals?
- Where is the total number of cases (n)?

Rates of Participation in Voluntary Associations by Marital Status for 100 Senior Citizens

| | Marita | | | |
|---------------------|---------------------------|-----------|---|------------------|
| Participation Rates | Married | Unmarried | | TOTALS |
| High | | | 1 | 50 |
| Low | parama rasp rassas con ya | | | <u>50</u> 100 |
| TOTALS | 50 | 50 | | 100 |



Important information to report

- Must have a title
- Cells are intersections of columns and rows
- Subtotals are called marginals
- Sample size (n) or population size (N) is reported at the intersection of row and column marginals



Independent, dependent variables

- Columns are scores of the independent variable
 - There will be as many columns as there are scores on the independent variable
- Rows are scores on the dependent variable
 - There will be as many rows as there are scores on the dependent variable
- Each cell reports the number of times each combination of scores occurred
 - There will be as many cells as there are scores on the two variables combined

Test for independence

- Chi square as a test of statistical significance is a test for independence
 - Two variables are independent if the classification of a case into a particular category of one variable has no effect on the probability that the case will fall into any particular category of the second variable

Rates of Participation in Voluntary Associations by Marital Status for 100 Senior Citizens

| | Marit | | |
|---------------------|---------|-----------------|-----------|
| Participation Rates | Married | Unmarried | TOTALS |
| High | 25 | 25 | 50 |
| Low | 25 | 25 | 50 |
| TOTALS | 50 | <u>25</u> 50 | 50 100 |

Cross tabulations

- Chi square is a test of significance based on bivariate tables
 - Bivariate tables are also called cross tabulations, crosstabs, contingency tables
- We are looking for significant differences between
 - The actual cell frequencies observed in a table (f_o)
 - And frequencies that would be expected by random chance or if cell frequencies were independent (f_e)



Computation of chi square

$$f_e = \frac{Row\ marginal \times Column\ marginal}{n}$$

$$\chi^2(obtained) = \sum \frac{(f_o - f_e)^2}{f_e}$$

where f_o = cell frequencies observed in the bivariate table

 f_e = cell frequencies that would be expected if the variables were independent

Example

- Random sample of 100 social work majors
 - We know whether the Council on Social Work Education has accredited their undergraduate programs
 - And whether they were hired in social work positions within three months of graduation
- Is there a significant relationship between employment status and accreditation status?

Employment of 100 Social Work Majors by Accreditation Status of Undergraduate Program

| | Accredi | | |
|--------------------------------|------------|----------------|--------|
| Employment Status | Accredited | Not Accredited | TOTALS |
| Working as a social worker | 30 | 10 | 40 |
| Not working as a social worker | 25 | 35 | 60 |
| TOTALS | 55 | 45 | 100 |

Step 1: Assumptions, requirements

- Independent random samples
- Level of measurement is nominal

- Note the minimal assumptions
 - No assumption is made about the shape of the sampling distribution
 - The chi square test is nonparametric or distributionfree



Step 2: Null hypothesis

- Null hypothesis, H_0 : $f_o = f_e$
 - The variables are independent
 - The observed frequencies are similar to the expected frequencies
- Alternative hypothesis, H_1 : $f_o \neq f_e$
 - The variables are dependent of each other
 - The observed frequencies are different than the expected frequencies



Step 3: Distribution, critical region

- Sampling distribution
 - Chi square distribution (χ^2)
- Significance level (α) = 0.05
 - The decision to reject the null hypothesis has only a 0.05 probability of being incorrect
- Degrees of freedom (df) = (r-1)(c-1)
 - -r = number of rows; c = number of columns
 - df = (r-1)(c-1) = (2-1)(2-1) = 1
- $\chi^2(critical) = 3.841$
 - If the probability (p-value) is less than 0.05
 - $-\chi^2(obtained)$ will be beyond $\chi^2(critical)$



Step 4: Test statistic

Observed frequencies

| | Accredi | | |
|--------------------------------|------------|----------------|--------|
| Employment Status | Accredited | Not Accredited | TOTALS |
| Working as a social worker | 30 | 10 | 40 |
| Not working as a social worker | 25 | 35 | 60 |
| TOTALS | 55 | 45 | 100 |

Expected frequencies

| | Accredit | | |
|--------------------------------|-------------|----------------|--------|
| Employment Status | Accredited | Not Accredited | TOTALS |
| Working as a social worker | 22 | 18 | 40 |
| Not working as a social worker | 33 | 27 | _60 |
| TOTALS | | 45 | 100 |

Expected frequency (f_e) for the top-left cell

$$f_e = \frac{Row\ marginal \times Column\ marginal}{n} = \frac{40 \times 55}{100} = 22$$

Computational table

| (1) | (2) | (3) | (4) | (5) |
|-------|------------|-----------------|----------------------------|---------------------|
| f_o | $f_{ m e}$ | $f_{o} - f_{e}$ | $(f_o - f_e)^2$ | $(f_o - f_e)^2/f_e$ |
| 30 | 22 | 8 | 64 | 2.91 |
| 10 | 18 | -8 | 64 | 3.56 |
| 25 | 33 | -8 | 64 | 1.94 |
| 35 | 27 | 8 | 64 | 2.37 |
| 100 | 100 | 0 | or all other cells can the | 10.78 |

•
$$\chi^2(obtained) = 10.78$$



Step 5: Decision, interpret

- $\chi^2(obtained) = 10.78$
 - This is beyond $\chi^2(critical) = 3.841$
 - The obtained χ^2 score falls in the critical region, so we reject the H_0
 - Therefore, the H₀ is false and must be rejected
- There is a significant relationship between employment status and accreditation status in the population from which the sample was drawn



Interpreting chi square

- The chi square test tells us only if the variables are independent or not
- It does not tell us the pattern or nature of the relationship
- To investigate the pattern, compute percentages within each column and compare across the columns



Limitations of chi square

- Difficult to interpret
 - When variables have many categories
 - Best when variables have four or fewer categories
- With small sample size (n)
 - We cannot assume that chi square sampling distribution will be accurate
 - Small samples: High percentage of cells have expected frequencies of 5 or less
- Like all tests of hypotheses
 - Chi square is sensitive to sample size
 - As n increases, obtained chi square increases
 - Large samples: Trivial relationships may be significant
- Statistical significance is not the same as substantive significance (importance, magnitude)



GSS example

- Is opinion about immigration different by sex?
- The probability of not rejecting H₀ is big (p>0.05)
 - Opinion about immigration does not depend on respondent's sex

. tab letin1 sex if year==2016, chi col

frequency
column percentage

| number of immigrants | rospondo | nts sov | |
|-----------------------|----------|----------|--------|
| to america nowadays | responde | IILS SEX | |
| should be | male | female | Total |
| increased a lot | 49 | 59 | 108 |
| | 5.98 | 5.75 | 5.85 |
| increased a little | 104 | 114 | 218 |
| | 12.70 | 11.11 | 11.82 |
| remain the same as it | 329 | 413 | 742 |
| | 40.17 | 40.25 | 40.22 |
| reduced a little | 181 | 238 | 419 |
| | 22.10 | 23.20 | 22.71 |
| reduced a lot | 156 | 202 | 358 |
| | 19.05 | 19.69 | 19.40 |
| Total | 819 | 1,026 | 1,845 |
| | 100.00 | 100.00 | 100.00 |

Source: 2016 General Social Survey.

Pearson chi2(4) = 1.3515

Pr = **0.853**

Edited table

Table 1. Opinion of the U.S. adult population about how should the number of immigrants to the country be nowadays by sex, 2004, 2010, and 2016

| Opinion About Number of Immigrants | Male (%) | Female (%) | Total (%) | Chi Square (df = 4) | p-value |
|---------------------------------------|-------------|---------------|--------------|------------------------|---------|
| 2004 | | | | 2.3397 | 0.6740 |
| Increase a lot | 3.17 | 4.30 | 3.78 | | |
| Increase a little | 6.89 | 6.27 | 6.56 | | |
| Remain the same | 35.01 | 34.05 | 34.49 | | |
| Reduce a little | 27.68 | 28.72 | 28.24 | | |
| Reduce a lot | 27.24 | 26.66 | 26.93 | | |
| Total | 100.00 | 100.00 | 100.00 | | |
| (sample size) | (914) | (1,069) | (1,983) | | |
| 2010 | | | | 7.0998 | 0.1310 |
| Increase a lot | 5.21 | 3.88 | 4.45 | | |
| Increase a little | 7.90 | 11.40 | 9.91 | | |
| Remain the same | 35.29 | 34.96 | 35.10 | | |
| Reduce a little | 24.03 | 25.31 | 24.77 | | |
| Reduce a lot | 27.56 | 24.44 | 25.77 | | |
| Total | 100.00 | 100.00 | 100.00 | | |
| (sample size) | (595) | (798) | (1,393) | | |
| 2016 | | | | 1.3515 | 0.8530 |
| Increase a lot | 5.98 | 5.75 | 5.85 | | |
| Increase a little | 12.70 | 11.11 | 11.82 | | |
| Remain the same | 40.17 | 40.25 | 40.22 | | |
| Reduce a little | 22.10 | 23.20 | 22.71 | | |
| Reduce a lot | 19.05 | 19.69 | 19.40 | | |
| Total | 100.00 | 100.00 | 100.00 | | |
| (sample size) | (819) | (1,026) | (1,845) | | |

Source: 2004, 2010, 2016 General Social Surveys.

ACS example

- Does education attainment vary by race/ethnicity?
 - The probability of not rejecting H_0 is small (p<0.01)
 - Education attainment is dependent on race/ethnicity
 - . tab educgr raceth [fweight=perwt], col nofreq

| | | | race | th | | | |
|-----------------------|--------|-----------|----------|--------|-----------|-----------|--------|
| educgr | White | African A | Hispanic | Asian | Native Am | Ohter rac | Total |
| Less than high school | 23.19 | 30.14 | 49.76 | 27.23 | 20.66 | 47.04 | 35.24 |
| High school | 26.55 | 29.72 | 26.11 | 16.23 | 34.00 | 17.85 | 26.09 |
| Some college | 20.38 | 22.79 | 14.40 | 12.29 | 25.15 | 16.42 | 17.82 |
| College | 19.92 | 11.04 | 7.12 | 23.26 | 15.36 | 12.51 | 13.78 |
| Graduate school | 9.95 | 6.31 | 2.62 | 20.99 | 4.83 | 6.17 | 7.07 |
| Total | 100.00 | 100.00 | 100.00 | 100.00 | 100.00 | 100.00 | 100.00 |

. svy: tab educgr raceth, col

(running tabulate on estimation sample)

Number of strata = 212 Number of PSUs = 114,016 Number of obs = 272,776 Population size = 28,995,881 Design df = 113,804

Pearson:

Uncorrected chi2(20) = 3.03e+04Design-based F(19.11, 2.2e+06)= 676.9183

P = 0.0000



Edited table

Table 1. Percentage distribution of population by educational attainment and race/ethnicity, Texas, 2019

| Educational attainment | Non- Hispanic White | Non- Hispanic Black | Hispanic | Non- Hispanic Asian | Non- Hispanic Native American | Other races | Total |
|-------------------------------------------|---------------------------|---------------------------|------------|---------------------------|----------------------------------------|-------------|------------|
| Less than high school | 23.19 | 30.14 | 49.76 | 27.23 | 20.66 | 47.04 | 35.24 |
| High school | 26.55 | 29.72 | 26.11 | 16.23 | 34.00 | 17.85 | 26.09 |
| Some college | 20.38 | 22.79 | 14.40 | 12.29 | 25.15 | 16.42 | 17.82 |
| College | 19.92 | 11.04 | 7.12 | 23.26 | 15.36 | 12.51 | 13.78 |
| Graduate school | 9.95 | 6.31 | 2.62 | 20.99 | 4.83 | 6.17 | 7.07 |
| Total | 100.00 | 100.00 | 100.00 | 100.00 | 100.00 | 100.00 | 100.00 |
| Population size (<i>N</i>) | 11,929,840 | 3,445,104 | 11,527,412 | 1,444,220 | 79,394 | 569,911 | 28,995,881 |
| Chi square (df = 20) | 3.03e+04 | | | | | | |
| Design-based <i>F</i> (19.11, 2.2e+06) | 676.92 | | | | | | |
| <i>p</i> -value | 0.0000 | | | | | | |

Source: 2019 American Community Survey.



